

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 December 2003 (04.12.2003)

PCT

(10) International Publication Number
WO 03/100008 A2

- (51) International Patent Classification⁷: C12N
- (21) International Application Number: PCT/US03/16283
- (22) International Filing Date: 22 May 2003 (22.05.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/383,459 24 May 2002 (24.05.2002) US
60/393,214 2 July 2002 (02.07.2002) US
60/436,254 23 December 2002 (23.12.2002) US
- (71) Applicant (for all designated States except US): **SCHERING CORPORATION** [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **WANG, Yan** [US/US]; 73 Clydesdale Road, Scotch Plains, NJ 07076 (US). **GREENBERG, Robert** [US/US]; 43 Whippoorwill Lane, Sparta, NJ 07871 (US). **PRESTA, Leonard** [US/US]; 1900 Gough Street, Apt. 206, San Francisco, CA 94109 (US). **PACHTER, Jonathan, A.** [US/US]; 13 Arcularius Terrace, Maplewood, NJ 07040 (US). **HAILEY, Judith** [US/US]; 84 Horizon Drive, Edison, NJ 08817 (US). **BRAMS, Peter** [DK/US]; 2740 11th Avenue, Sacramento, CA 95818 (US). **WILLIAMS, Denise** [US/US]; 5511 Chesbro Avenue, San Jose, CA 95123 (US). **SRINIVASAN, Mohan** [IN/US]; 21073 Red Fir Court, Cupertino, CA 95014 (US). **FEINGERSH, Diane** [US/US]; 204 Appian Way, Union City, CA 94587 (US).
- (81) Designated States (national): AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV, MA, MD, MG, MK, MN, MX, MZ, NI, NO, NZ, PH, PL, PT, RO, RU, SC, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, US, UZ, VC, VN, YU, ZA, ZM.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declarations under Rule 4.17:**
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
 - as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
 - as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- Published:**
- without international search report and to be republished upon receipt of that report
 - with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/100008 A2

(54) Title: NEUTRALIZING HUMAN ANTI-IGFR ANTIBODY

(57) Abstract: The present invention includes fully human, neutralizing, monoclonal antibodies against human Insulin-like Growth Factor Receptor-I (IGFR1). The antibodies are useful for treating or preventing cancer in a subject. Also included are methods of using and producing the antibodies of the invention.

NEUTRALIZING HUMAN ANTI-IGFR ANTIBODY

This application claims the benefit of U.S. Provisional Patent Application No. 60/383,459, filed May 24, 2002; U.S. Provisional Patent Application No. 60/393,214, filed July 2, 2002 and U.S. Provisional Patent Application No. 60/436,254, filed December 23, 2002 each of which is herein incorporated by reference in its entirety.

FIELD OF THE INVENTION

The present invention relates to fully human, monoclonal anti-Insulin-like Growth Factor Receptor-I (IGFR1) antibodies as well as methods of using the antibodies and methods of producing the antibodies.

5

BACKGROUND OF THE INVENTION

The insulin-like growth factors, also known as somatomedins, include insulin-like growth factor-I (IGF-I) and insulin-like growth factor-II (IGF-II) (Klapper, *et al.*, (1983) *Endocrinol.* 112:2215 and Rinderknecht, *et al.*, (1978) *Febs.Lett.* 89:283). These growth factors exert mitogenic activity on various cell types, including tumor cells (Macaulay, (1992) *Br. J. Cancer* 65:311), by binding to a common receptor named the insulin-like growth factor receptor-1 (IGFR1) (Sepp-Lorenzino, (1998) *Breast Cancer Research and Treatment* 47:235). Interaction of IGFs with IGFR1 activates the receptor by triggering autophosphorylation of the receptor on tyrosine residues (Butler, *et al.*, (1998) *Comparative Biochemistry and Physiology* 121:19).
10 Once activated, IGFR1, in turn, phosphorylates intracellular targets to activate cellular signaling pathways. This receptor activation is critical for stimulation of tumor cell growth and survival. Therefore, inhibition of IGFR1 activity represents a valuable potential method to treat or prevent growth of human cancers and other proliferative diseases.

20

Several lines of evidence indicate that IGF-I, IGF-II and their receptor IGFR1 are important mediators of the malignant phenotype. Plasma levels of IGF-I have been found to be the strongest predictor of prostate cancer risk (Chan, *et al.*, (1998) *Science* 279:563) and similar epidemiological studies strongly link plasma IGF-I levels with breast, colon and lung cancer risk.

Overexpression of Insulin-like Growth Factor Receptor-I has also been demonstrated in several cancer cell lines and tumor tissues. IGFR1 is overexpressed in 40% of all breast cancer cell lines (Pandini, *et al.*, (1999) Cancer Res. 5:1935) and in 15% of lung cancer cell lines. In breast cancer tumor tissue, IGFR1 is
5 overexpressed 6-14 fold and IGFR1 exhibits 2-4 fold higher kinase activity as compared to normal tissue (Webster, *et al.*, (1996) Cancer Res. 56:2781 and Pekonen, *et al.*, (1998) Cancer Res. 48:1343). Ninety percent of colorectal cancer tissue biopsies exhibit elevated IGFR1 levels wherein the extent of IGFR1 expression is correlated with the severity of the disease. Analysis of primary cervical cancer cell
10 cultures and cervical cancer cell lines revealed 3- and 5-fold overexpression of IGFR1, respectively, as compared to normal ectocervical cells (Steller, *et al.*, (1996) Cancer Res. 56:1762). Expression of IGFR1 in synovial sarcoma cells also correlated with an aggressive phenotype (*i.e.*, metastasis and high rate of proliferation; Xie, *et al.*, (1999) Cancer Res. 59:3588).

15 Acromegaly, a slowly developing disease, is caused by hypersecretion of growth hormone and IGF-I (Ben-Schlomo, *et al.*, (2001) Endocrin. Metab.Clin. North. Am. 30:565-583). Antagonism of IGFR1 function may be helpful in treating the disease.

There are several antibodies, which are known in the art, which inhibit the
20 activity of IGFR1. However, these are of relatively low therapeutic value. For example, α -IR3 (Kull, *et al.*, (1983) J. Biol. Chem. 258:6561), 1H7 (Li *et al.*, (1993) Biochem. Biophys. Res. Comm. 196:92-98 and Xiong *et al.*, (1992) Proc. Natl. Acad. Sci., U.S.A. 89:5356-5360; Santa Cruz biotechnology, Inc.; Santa Cruz, CA) and MAB391 (R&D Systems; Minneapolis, MN) are mouse monoclonal antibodies which
25 interact with IGFR1 and inhibit its activity. Since these are mouse antibodies, their therapeutic utility in humans is limited. When immunocompetent human subjects are administered a dose of mouse antibodies, the subjects produce antibodies against the mouse immunoglobulin sequences. These human anti-mouse antibodies (HAMA) neutralize the therapeutic antibodies and may induce acute toxicity (*i.e.*, a HAMA
30 response).

One method by which to avert a HAMA response is through the use of fully-human antibodies which lack any foreign (*e.g.*, mouse) amino acid sequences. Although the use of fully-human antibodies is an effective method by which to reduce or prevent human host immune rejection of the therapeutic antibody, rejection of the

fully-human antibody can occur. Human rejection of human antibodies may be referred to as a human anti-human antibody response (HAHA response). HAHA response can be mediated by factors such as the presence of rare, low occurrence amino acid sequences in the fully-human antibodies. For this reason, therapeutic antibodies may also be optimized by the inclusion of non-immunogenic or only weakly immunogenic human antibody framework sequences. Preferably, the sequences occur frequently in other human antibodies.

SUMMARY OF THE INVENTION

10 The present invention provides fully human anti-human IGFR1 monoclonal antibodies which, preferably, will not induce a HAMA response or will not induce a HAHA response when administered to human subjects and which are useful for treating or preventing diseases which are mediated by IGFR1 (e.g., malignancy).

15 The present invention provides a binding composition (e.g., an antibody or antigen-binding fragment thereof) comprising a light chain, wherein the chain comprises the amino acid sequence of the light chain CDR-L1 defined by SEQ ID NO: 8 or 31, the amino acid sequence of the light chain CDR-L2 defined by SEQ ID NO: 9 or 32 and the amino acid sequence of the light chain CDR-L3 defined by SEQ ID NO: 10 or 33. Also provided is a binding composition (e.g., an antibody or antigen-binding fragment thereof) including a heavy chain, wherein the chain includes the amino acid sequence of the heavy chain CDR-H1 defined by SEQ ID NO: 14 or 37, the amino acid sequence of the heavy chain CDR-H2 defined by SEQ ID NO: 15 or 38 and the amino acid sequence of the heavy chain CDR-H3 defined by SEQ ID NO: 16 or 39.

25 Preferably, the binding composition (e.g., an antibody or antigen-binding fragment thereof) of the invention comprises a light chain variable region, preferably a mature light chain variable region, which includes amino acids 20-128 of SEQ ID NO: 2, amino acids 21-130 of SEQ ID NO: 25, amino acids 20-128 of SEQ ID NO: 41 or 43 or amino acids 20-128 of SEQ ID NO: 41, 43, 72, 74, 76 or 78 and/or a heavy chain variable region, preferably a mature heavy chain variable region, which includes amino acids 20-137 of SEQ ID NO: 4, amino acids 20-140 of SEQ ID NO: 27, amino acids 20-137 of SEQ ID NO: 45 or amino acids 20-137 of SEQ ID NO: 112.

30 Pharmaceutical compositions comprising a binding composition of the present invention and a pharmaceutically acceptable carrier are also provided by the present

invention. The binding composition of the invention may also be conjugated to a substance such as polyethylene glycol.

The present invention also includes a binding composition (*e.g.*, a human antibody or antigen binding fragment thereof) which specifically binds to human

- 5 IGFR1 comprising a property selected from the group consisting of:
- (a) binds to IGFR1 (*e.g.*, human IGFR1) with a K_d of about 86×10^{-11} or less;
 - (b) Has an off rate (K_{off}) for IGFR1 (*e.g.*, human IGFR1) of about 6.50×10^{-5} or smaller;
 - (c) Has an on rate (K_{on}) for IGFR1 (*e.g.*, human IGFR1) of about 0.7×10^5 or greater;
 - 10 (d) Competes with IGF1 for binding to IGFR1 (*e.g.*, human IGFR1);
 - (e) Inhibits autophosphorylation (*e.g.*, with an IC_{50} of 0.10 nM) of IGFR1 (*e.g.*, human IGFR1); and
 - (f) Inhibits anchorage-independent growth of a cell expressing IGFR1 (*e.g.*, human IGFR1).
- 15 Preferably, the binding composition comprises all of said properties (a-f). More preferably, the binding composition (*e.g.*, a human antibody or antigen binding fragment thereof) comprises a member selected from:
- (a) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 8, CDR-L2 defined by SEQ ID NO: 9 and CDR-L3 defined by SEQ ID NO: 10;
 - 20 (b) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 31, CDR-L2 defined by SEQ ID NO: 32 and CDR-L3 defined by SEQ ID NO: 33;
 - (c) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 14 or SEQ ID NO: 17, CDR-H2 defined by SEQ ID NO: 15 and CDR-H3 defined by SEQ ID NO: 16; and
 - 25 (d) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 37 or SEQ ID NO: 70, CDR-H2 defined by SEQ ID NO: 38 and CDR-H3 defined by SEQ ID NO: 39.

The present invention also includes an isolated nucleic acid encoding a

30 peptide selected from:

- (a) amino acids 20-128 of SEQ ID NO: 2;
- (b) amino acids 21-130 of SEQ ID NO: 25;
- (c) amino acids 20-128 of SEQ ID NO: 72;
- (d) amino acids 20-128 of SEQ ID NO: 74;

- (a) amino acids 20-137 of SEQ ID NO: 4;
- (b) amino acids 20-140 of SEQ ID NO: 27;
- (c) amino acids 20-137 of SEQ ID NO: 45;
- (d) amino acids 20-137 of SEQ ID NO: 112;
- 5 (e) amino acids 20-128 of SEQ ID NO: 76; and
- (f) amino acids 20-128 of SEQ ID NO: 78.

Preferably, the nucleic acid is selected from:

- (a) nucleotides 58-384 of SEQ ID NO: 1;
- (b) nucleotides 61-390 of SEQ ID NO: 24;
- 10 (c) nucleotides 58-384 of SEQ ID NO: 71;
- (d) nucleotides 58-384 of SEQ ID NO: 73.
- (e) nucleotides 58-411 of SEQ ID NO: 3;
- (f) nucleotides 58-420 of SEQ ID NO: 26;
- (g) nucleotides 58-411 of SEQ ID NO: 44;
- 15 (h) nucleotides 58-411 of SEQ ID NO: 111;
- (i) nucleotides 58-384 of SEQ ID NO: 75; and
- (j) nucleotides 58-384 of SEQ ID NO: 77.

The present invention also provides a recombinant vector comprising any of the foregoing polynucleotides along with a host cell comprising the vector.

- 20 The present invention also comprises a polypeptide selected from:

- (a) amino acids 20-128 of SEQ ID NO: 2;
- (b) amino acids 21-130 of SEQ ID NO: 25;
- (c) amino acids 20-128 of SEQ ID NO: 72;
- (d) amino acids 20-128 of SEQ ID NO: 74;
- 25 (e) amino acids 20-137 of SEQ ID NO: 4;
- (f) amino acids 20-140 of SEQ ID NO: 27;
- (g) amino acids 20-137 of SEQ ID NO: 45;
- (h) amino acids 20-137 of SEQ ID NO: 112;
- (i) amino acids 20-128 of SEQ ID NO: 76; and
- 30 (j) amino acids 20-128 of SEQ ID NO: 78.

Preferably, the binding composition of the present invention is a human antibody comprising at least one (*e.g.*, 1 or 2) light chain/heavy chain combination selected from:

- a) a light chain variable region comprising amino acids 20-128 of SEQ ID NO: 2 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 4; [15H12/19D12 mature LC – 15H12/19D12 mature HC]
- b) a light chain variable region comprising amino acids 21-130 of SEQ ID NO: 25 and a heavy chain variable region comprising amino acids 20-140 of SEQ ID NO: 27; [1H3 mature LC – 1H3 mature HC]
- c) a light variable region comprising amino acids 20-128 of SEQ ID NO: 72 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 45; [mature LCC - mature HCA]
- d) a light variable region comprising amino acids 20-128 of SEQ ID NO: 74 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 45; [mature LCD - mature HCA]
- e) a light variable region comprising amino acids 20-128 of SEQ ID NO: 76 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 45; [mature LCE - mature HCA]
- f) a light variable region comprising amino acids 20-128 of SEQ ID NO: 78 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 45; [mature LCF - mature HCA]
- g) a light variable region comprising amino acids 20-128 of SEQ ID NO: 72 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 112; [mature LCC - mature HCB]
- h) a light variable region comprising amino acids 20-128 of SEQ ID NO: 74 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 112; [mature LCD - mature HCB]
- i) a light variable region comprising amino acids 20-128 of SEQ ID NO: 76 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 112; [mature LCE - mature HCB] and
- j) a light variable region comprising amino acids 20-128 of SEQ ID NO: 78 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 112. [mature LCF - mature HCB].

More preferably, the human antibody is a tetramer comprising two of the foregoing light/heavy chain pairs. Preferably, the human antibody includes mature LCF paired with mature HCA or mature HCB.

Also provided is a method for making a polypeptide comprising amino acids 20-128 of SEQ ID NO: 2, amino acids 20-137 of SEQ ID NO: 4, amino acids 21-130 of SEQ ID NO: 25, amino acids 20-140 of SEQ ID NO: 27, amino acids 20-128 of SEQ ID NO: 41, 43, 72, 74, 76 or 78, amino acids 20-137 of SEQ ID NO: 45 or amino acids 20-137 of SEQ ID NO: 112 comprising culturing the host cell under conditions in which the polypeptide is produced. Preferably, the polypeptide is also isolated from the host cell.

The invention also provides a method for treating or preventing a medical condition in a subject which is mediated by elevated expression or activity of Insulin-like Growth Factor Receptor-I or by elevated expression of one or more of its ligands (*e.g.*, IGF-I or IGF-II) comprising administering a binding composition of the invention (*e.g.*, antibody or antigen-binding fragment of the invention) to the subject. Preferably, the binding composition comprises a member selected from:

- (a) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 8, CDR-L2 defined by SEQ ID NO: 9 and CDR-L3 defined by SEQ ID NO: 10;
- (b) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 31, CDR-L2 defined by SEQ ID NO: 32 and CDR-L3 defined by SEQ ID NO: 33;
- (c) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 14 or SEQ ID NO: 17, CDR-H2 defined by SEQ ID NO: 15 and CDR-H3 defined by SEQ ID NO: 16; and
- (d) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 37 or SEQ ID NO: 70, CDR-H2 defined by SEQ ID NO: 38 and CDR-H3 defined by SEQ ID NO: 39.

The present invention includes any plasmid selected from the group consisting of :

- (i) CMV promoter-15H12/19D12 HCA (γ 4)-
Deposit name: "15H12/19D12 HCA (γ 4)";
ATCC accession No.: _____;
- (ii) CMV promoter-15H12/19D12 HCB (γ 4)-
Deposit name: "15H12/19D12 HCB (γ 4)";
ATCC accession No.: _____;
- (iii) CMV promoter-15H12/19D12 HCA (γ 1)-
Deposit name: "15H12/19D12 HCA (γ 1)";

ATCC accession No.: _____;

(iv) CMV promoter-15H12/19D12 LCC (κ)-

Deposit name: "15H12/19D12 LCC (κ)";

ATCC accession No.: _____;

5 (v) CMV promoter-15H12/19D12 LCD (κ)-

Deposit name: "15H12/19D12 LCD (κ)";

ATCC accession No.: _____;

(vi) CMV promoter-15H12/19D12 LCE (κ)-

Deposit name: "15H12/19D12 LCE (κ)";

10 ATCC accession No.: _____; and

(vii) CMV promoter-15H12/19D12 LCF (κ)-

Deposit name: "15H12/19D12 LCF (κ)";

ATCC accession No.: _____;

as well as the nucleic acid inserts of any of the foregoing plasmids. Also included are
15 the nucleic acid portions of the inserts encoding the immunoglobulin variable regions
included in the plasmid inserts optionally including the immunoglobulin constant
region (*i.e.*, excluding the signal sequence). Also included are any polypeptides
encoded by the nucleic acids of any of the foregoing plasmid inserts as well as
polypeptides encoding the immunoglobulin variable regions included in any insert
20 optionally including the immunoglobulin constant region (*i.e.*, excluding the signal
sequence).

The above-identified plasmids were deposited, under the Budapest Treaty, on
_____ with the American Type Culture Collection (ATCC); 10801 University
Boulevard; Manassas, Virginia 20110-2209. All restrictions on access to the plasmids
25 deposited in ATCC will be removed upon grant of a patent.

Preferably, the binding composition is combined with a pharmaceutically
acceptable carrier in a pharmaceutical composition. Such medical conditions, as
contemplated by the present invention, include acromegaly, ovarian cancer,
pancreatic cancer, benign prostatic hyperplasia, breast cancer, prostate cancer, bone
30 cancer, lung cancer, colorectal cancer, cervical cancer, synovial sarcoma, diarrhea
associated with metastatic carcinoid, vasoactive intestinal peptide secreting tumors,
gigantism, psoriasis, atherosclerosis, smooth muscle restenosis of blood vessels and
inappropriate microvascular proliferation.

The binding compositions may be administered to a subject, for example, by a parenteral route. Combination therapies comprising administration of a binding composition of the present invention in association with an anti-cancer therapy agent or in association with an anti-cancer therapeutic procedure are also provided.

- 5 A method for producing a fully-human anti-IGFR1 antibody which comprises the steps of immunizing a transgenic non-human animal having a genome comprising a human heavy chain transgene and a human light chain transgene with IGFR1 antigenic polypeptide, preferably amino acids 30-902 of SEQ ID NO: 19 and/or a cell (e.g., HEK293) which expresses IGFR1 on its surface, such that antibodies are
10 produced by B cells of the animal; isolating B cells of the animal; fusing the B cells with myeloma cells to form immortal, hybridoma cells that secrete human monoclonal antibodies specific for IGFR1; and isolating the human monoclonal antibodies specific for IGFR1 is also provided.

15

DETAILED DESCRIPTION

- Preferred embodiments of the present invention include a fully human, monoclonal antibody or antigen-binding fragment thereof which specifically recognizes and binds to Insulin-like Growth Factor Receptor-I, preferably amino acids 30-902 of SEQ ID NO: 19. Preferably, the antibody or antigen-binding fragment
20 thereof is 1H3, 15H12, 19D12, 15H12/19D12 LCA, 15H12/19D12 LCB, 15H12/19D12 LCC, 15H12/19D12 LCD, 15H12/19D12 LCE, 15H12/19D12 LCF, 15H12/19D12 HCA or 15H12/19D12 HCB.

- A binding composition or agent refers to a molecule that binds with specificity to IGFR1, e.g., in a ligand-receptor type fashion or an antibody-antigen interaction,
25 e.g., proteins which specifically associate with IGFR1, e.g., in a natural physiologically relevant protein-protein interaction, either covalent or non-covalent. The term "binding composition" is preferably a polypeptide, such as a full antibody or antigen-binding fragment thereof of the present invention (e.g., 15H12/19D12 LCA, 15H12/19D12 LCB, 15H12/19D12 LCC, 15H12/19D12 LCD, 15H12/19D12 LCE,
30 15H12/19D12 LCF, 15H12/19D12 HCA OR 15H12/19D12 HCB or any peptide set forth, below, in Table 1).

The antibodies and antigen-binding fragments of the invention may be used to inhibit growth of cells, preferably malignant cells, both *in vitro* and *in vivo*. Without being bound by a single theory, the antibodies and antigen-binding fragments of the

invention may inhibit cellular growth by inhibiting the interaction between IGFR1 and a ligand for the receptor, such as Insulin-like Growth Factor-I (IGF-I) or Insulin-like Growth Factor-II (IGF-II). The antibodies and antigen-binding fragments may also inhibit IGFR1 autophosphorylation, inhibit anchorage-independent growth of cells
5 (e.g., cancer cells) expressing IGFR1 and inhibit activation of AKT kinase by inducing degradation of IGFR1. Preferably, the antibodies and antigen-binding fragments neutralize the activity of IGFR1 and/or down-regulate IGFR1. The antibodies and antigen-binding fragments may be used to treat or prevent diseases which are mediated by IGFR1. The present invention also provides methods for making the
10 antibodies and antigen-binding fragments of the invention.

The term "antibody molecule" refers to whole antibodies (e.g., IgG, preferably, IgG1 or IgG4) and fragments, preferably antigen-binding fragments, thereof. Antibody fragments include Fab antibody fragments, F(ab)₂ antibody fragments, Fv antibody fragments, single chain Fv antibody fragments and dsFv antibody fragments.

15 The terms "IGFR1" "Insulin-like Growth Factor Receptor-I" and "Insulin-like Growth Factor Receptor, type I" are well known in the art. Although IGFR1 may be from any organism, it is preferably from an animal, more preferably from a mammal (e.g., mouse, rat, rabbit, sheep or dog) and most preferably from a human. The nucleotide and amino acid sequence of a typical human IGFR1 precursor has the
20 Genbank Accession No. X04434 or NM_000875 (SEQ ID NO: 19). Cleavage of the precursor (e.g., between amino acids 710 and 711) produces an α -subunit and a β -subunit which associate to form a mature receptor. In preferred embodiments of the invention, amino acids 30-902, from the full length IGFR1 polypeptide are used as an antigen for generation of anti-IGFR1 antibodies.

25 The terms "IGF-I" "Insulin-like Growth Factor-I" and "Insulin-like Growth Factor, type I" are also well known in the art. The terms "IGF-II" "Insulin-like Growth Factor-II" and "Insulin-like Growth Factor, type II" are also well known in the art. Although IGF-I or IGF-II may be from any organism, they are preferably from an animal, more preferably from a mammal (e.g., mouse, rat, rabbit, sheep or dog) and
30 most preferably from a human. The nucleic acid and amino acid sequence of typical, human IGF-I and IGF-II have the Genbank Accession No. XM_052648 (SEQ ID NO: 20) and NM_000612 (SEQ ID NO: 21), respectively. The term "sIGFR1" or "soluble IGFR1" includes any soluble fragment of IGFR1 (e.g., human IGFR1), preferably a

fragment from which the receptor trans-membrane region has been deleted, more preferably amino acids 30-902 of SEQ ID NO: 19.

The amino acid sequence of the variable region of preferred, fully human, monoclonal anti-IGFR1 antibody molecules of the invention (*e.g.*, 1H3, 15H12 and 19D12) along with the nucleotide sequences of nucleic acids which encode the regions are summarized in Table 1. The present invention includes any nucleic acid or polypeptide (*e.g.*, antibody) which comprises one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7 or 8) of any of the nucleic acids or polypeptides (including mature fragments thereof) set forth, below, in Table 1. Table 1 also includes a summary of the amino acid and nucleotide sequences which correspond to the CDR regions of the antibodies. The amino acid and nucleotide sequences corresponding to the variable region of 15H12 and 19D12 are identical; for this reason, only a single sequence for each variable region or CDR is shown.

Table 1. Summary of amino acid and nucleotide sequences of the invention.

SEQUENCE	SEQUENCE IDENTIFIER
Nucleotide sequence encoding the 15H12 and 19D12 light chain variable region-including signal peptide (15H12/19D12 LC)	SEQ ID NO: 1
Amino acid sequence of the 15H12 and 19D12 light chain variable region-including signal peptide	SEQ ID NO: 2
Nucleotide sequence encoding the 15H12 and 19D12 heavy chain variable region including signal peptide (15H12/19D12 HC)	SEQ ID NO: 3
Amino acid sequence of the 15H12 and 19D12 heavy chain variable region including signal peptide	SEQ ID NO: 4
Nucleotide sequence encoding the 15H12 and 19D12 CDR-L1	SEQ ID NO: 5
Nucleotide sequence encoding the 15H12 and 19D12 CDR-L2	SEQ ID NO: 6
Nucleotide sequence encoding the 15H12 and 19D12 CDR-L3	SEQ ID NO: 7
Amino acid sequence of the 15H12 and 19D12 CDR-L1	SEQ ID NO: 8

SEQUENCE	SEQUENCE IDENTIFIER
Amino acid sequence of the 15H12 and 19D12 CDR-L2	SEQ ID NO: 9
Amino acid sequence of the 15H12 and 19D12 CDR-L3	SEQ ID NO: 10
Nucleotide sequence encoding the 15H12 and 19D12 CDR-H1	SEQ ID NO: 11
Nucleotide sequence encoding the 15H12 and 19D12 CDR-H2	SEQ ID NO: 12
Nucleotide sequence encoding the 15H12 and 19D12 CDR-H3	SEQ ID NO: 13
Amino acid sequence of the 15H12 and 19D12 CDR-H1	SEQ ID NO: 14
Amino acid sequence of the 15H12 and 19D12 CDR-H2	SEQ ID NO: 15
Amino acid sequence of the 15H12 and 19D12 CDR-H3	SEQ ID NO: 16
Amino acid sequence of an alternative 15H12 and 19D12 CDR-H1	SEQ ID NO: 17
Nucleotide sequence encoding an alternative 15H12 and 19D12 CDR-H1	SEQ ID NO: 18
Amino acid sequence of Insulin-like Growth Factor Receptor-I (IGFR1)	SEQ ID NO: 19
Amino acid sequence of Insulin-like Growth Factor-I (IGF1)	SEQ ID NO: 20
Amino acid sequence of Insulin-like Growth Factor-II (IGF2)	SEQ ID NO: 21
Nucleotide sequence of PCR primer	SEQ ID NO: 22
Nucleotide sequence of PCR primer	SEQ ID NO: 23
Nucleotide sequence encoding the 1H3 light chain variable region-including signal peptide (1H3 LC)	SEQ ID NO: 24
Amino acid sequence of the 1H3 light chain variable region-including signal peptide	SEQ ID NO: 25
Nucleotide sequence encoding the 1H3 heavy chain variable region including signal peptide (1H3 HC)	SEQ ID NO: 26
Amino acid sequence of the 1H3 heavy chain variable region including signal peptide	SEQ ID NO: 27

SEQUENCE	SEQUENCE IDENTIFIER
Nucleotide sequence encoding the 1H3 CDR-L1	SEQ ID NO: 28
Nucleotide sequence encoding the 1H3 CDR-L2	SEQ ID NO: 29
Nucleotide sequence encoding the 1H3 CDR-L3	SEQ ID NO: 30
Amino acid sequence of the 1H3 CDR-L1	SEQ ID NO: 31
Amino acid sequence of the 1H3 CDR-L2	SEQ ID NO: 32
Amino acid sequence of the 1H3 CDR-L3	SEQ ID NO: 33
Nucleotide sequence encoding the 1H3 CDR-H1	SEQ ID NO: 34
Nucleotide sequence encoding the 1H3 CDR-H2	SEQ ID NO: 35
Nucleotide sequence encoding the 1H3 CDR-H3	SEQ ID NO: 36
Amino acid sequence of the 1H3 CDR-H1	SEQ ID NO: 37
Amino acid sequence of the 1H3 CDR-H2	SEQ ID NO: 38
Amino acid sequence of the 1H3 CDR-H3	SEQ ID NO: 39
Nucleotide sequence encoding the 15H12/19D12 light chain A (LCA)	SEQ ID NO:40
Amino acid sequence of the 15H12/19D12 light chain A	SEQ ID NO:41
Nucleotide sequence encoding the 15H12/19D12 light chain B (LCB)	SEQ ID NO:42
Amino acid sequence of the 15H12/19D12 light chain B	SEQ ID NO:43
Nucleotide sequence encoding the 15H12/19D12 heavy chain A (HCA)	SEQ ID NO:44
Amino acid sequence of the 15H12/19D12 heavy chain A	SEQ ID NO:45
Nucleotide sequence encoding the 15H12/19D12 light chain A framework region 1	SEQ ID NO:46
Amino acid sequence of the 15H12/19D12 light chain A framework region 1	SEQ ID NO:47
Nucleotide sequence encoding the 15H12/19D12 light chain A framework region 2	SEQ ID NO:48

SEQUENCE	SEQUENCE IDENTIFIER
Amino acid sequence of the 15H12/19D12 light chain A framework region 2	SEQ ID NO:49
Nucleotide sequence encoding the 15H12/19D12 light chain A framework region 3	SEQ ID NO:50
Amino acid sequence of the 15H12/19D12 light chain A framework region 3	SEQ ID NO:51
Nucleotide sequence encoding the 15H12/19D12 light chain A framework region 4	SEQ ID NO:52
Amino acid sequence of the 15H12/19D12 light chain A framework region 4	SEQ ID NO:53
Nucleotide sequence encoding the 15H12/19D12 light chain B framework region 1	SEQ ID NO:54
Amino acid sequence of the 15H12/19D12 light chain B framework region 1	SEQ ID NO:55
Nucleotide sequence encoding the 15H12/19D12 light chain B framework region 2	SEQ ID NO:56
Amino acid sequence of the 15H12/19D12 light chain B framework region 2	SEQ ID NO:57
Nucleotide sequence encoding the 15H12/19D12 light chain B framework region 3	SEQ ID NO:58
Amino acid sequence of the 15H12/19D12 light chain B framework region 3	SEQ ID NO:59
Nucleotide sequence encoding the 15H12/19D12 light chain B framework region 4	SEQ ID NO:60
Amino acid sequence of the 15H12/19D12 light chain B framework region 4	SEQ ID NO:61
Nucleotide sequence encoding the 15H12/19D12 heavy chain A framework region 1	SEQ ID NO:62
Amino acid sequence of the 15H12/19D12 heavy chain A framework region 1	SEQ ID NO:63
Nucleotide sequence encoding the 15H12/19D12 heavy chain A framework region 2	SEQ ID NO:64

SEQUENCE	SEQUENCE IDENTIFIER
Amino acid sequence of the 15H12/19D12 heavy chain A framework region 2	SEQ ID NO:65
Nucleotide sequence encoding the 15H12/19D12 heavy chain A framework region 3	SEQ ID NO:66
Amino acid sequence of the 15H12/19D12 heavy chain A framework region 3	SEQ ID NO:67
Nucleotide sequence encoding the 15H12/19D12 heavy chain A framework region 4	SEQ ID NO:68
Amino acid sequence of the 15H12/19D12 heavy chain A framework region 4	SEQ ID NO:69
Amino acid sequence of the alternative 1H3 CDR-H1	SEQ ID NO: 70
Nucleotide sequence encoding the 15H12/19D12 light chain C (LCC)	SEQ ID NO: 71
Amino acid sequence of the 15H12/19D12 light chain C	SEQ ID NO: 72
Nucleotide sequence encoding the 15H12/19D12 light chain D (LCD)	SEQ ID NO: 73
Amino acid sequence of the 15H12/19D12 light chain D	SEQ ID NO: 74
Nucleotide sequence encoding the 15H12/19D12 light chain E (LCE)	SEQ ID NO: 75
Amino acid sequence of the 15H12/19D12 light chain E	SEQ ID NO: 76
Nucleotide sequence encoding the 15H12/19D12 light chain F (LCF)	SEQ ID NO: 77
Amino acid sequence of the 15H12/19D12 light chain F	SEQ ID NO: 78
Nucleotide sequence encoding the 15H12/19D12 light chain C framework region 1	SEQ ID NO: 79
Amino acid sequence of the 15H12/19D12 light chain C framework region 1	SEQ ID NO: 80
Nucleotide sequence encoding the 15H12/19D12 light chain C framework region 2	SEQ ID NO: 81

SEQUENCE	SEQUENCE IDENTIFIER
Amino acid sequence of the 15H12/19D12 light chain C framework region 2	SEQ ID NO: 82
Nucleotide sequence encoding the 15H12/19D12 light chain C framework region 3	SEQ ID NO: 83
Amino acid sequence of the 15H12/19D12 light chain C framework region 3	SEQ ID NO: 84
Nucleotide sequence encoding the 15H12/19D12 light chain C framework region 4	SEQ ID NO: 85
Amino acid sequence of the 15H12/19D12 light chain C framework region 4	SEQ ID NO: 86
Nucleotide sequence encoding the 15H12/19D12 light chain D framework region 1	SEQ ID NO: 87
Amino acid sequence of the 15H12/19D12 light chain D framework region 1	SEQ ID NO: 88
Nucleotide sequence encoding the 15H12/19D12 light chain D framework region 2	SEQ ID NO: 89
Amino acid sequence of the 15H12/19D12 light chain D framework region 2	SEQ ID NO: 90
Nucleotide sequence encoding the 15H12/19D12 light chain D framework region 3	SEQ ID NO: 91
Amino acid sequence of the 15H12/19D12 light chain D framework region 3	SEQ ID NO: 92
Nucleotide sequence encoding the 15H12/19D12 light chain D framework region 4	SEQ ID NO: 93
Amino acid sequence of the 15H12/19D12 light chain D framework region 4	SEQ ID NO: 94
Nucleotide sequence encoding the 15H12/19D12 light chain E framework region 1	SEQ ID NO: 95
Amino acid sequence of the 15H12/19D12 light chain E framework region 1	SEQ ID NO: 96
Nucleotide sequence encoding the 15H12/19D12 light chain E framework region 2	SEQ ID NO: 97

SEQUENCE	SEQUENCE IDENTIFIER
Amino acid sequence of the 15H12/19D12 light chain E framework region 2	SEQ ID NO: 98
Nucleotide sequence encoding the 15H12/19D12 light chain E framework region 3	SEQ ID NO: 99
Amino acid sequence of the 15H12/19D12 light chain E framework region 3	SEQ ID NO: 100
Nucleotide sequence encoding the 15H12/19D12 light chain E framework region 4	SEQ ID NO: 101
Amino acid sequence of the 15H12/19D12 light chain E framework region 4	SEQ ID NO: 102
Nucleotide sequence encoding the 15H12/19D12 light chain F framework region 1	SEQ ID NO: 103
Amino acid sequence of the 15H12/19D12 light chain F framework region 1	SEQ ID NO: 104
Nucleotide sequence encoding the 15H12/19D12 light chain F framework region 2	SEQ ID NO: 105
Amino acid sequence of the 15H12/19D12 light chain F framework region 2	SEQ ID NO: 106
Nucleotide sequence encoding the 15H12/19D12 light chain F framework region 3	SEQ ID NO: 107
Amino acid sequence of the 15H12/19D12 light chain F framework region 3	SEQ ID NO: 108
Nucleotide sequence encoding the 15H12/19D12 light chain F framework region 4	SEQ ID NO: 109
Amino acid sequence of the 15H12/19D12 light chain F framework region 4	SEQ ID NO: 110
Nucleotide sequence encoding the 15H12/19D12 heavy chain B (HCB)	SEQ ID NO: 111
Amino acid sequence of the 15H12/19D12 heavy chain B	SEQ ID NO: 112
Nucleotide sequence encoding the 15H12/19D12 heavy chain B framework region 1	SEQ ID NO: 113

SEQUENCE	SEQUENCE IDENTIFIER
Amino acid sequence of the 15H12/19D12 heavy chain B framework region 1	SEQ ID NO: 114
Nucleotide sequence encoding the 15H12/19D12 heavy chain B F framework region 2	SEQ ID NO: 115
Amino acid sequence of the 15H12/19D12 heavy chain B framework region 2	SEQ ID NO: 116
Nucleotide sequence encoding the 15H12/19D12 heavy chain B framework region 3	SEQ ID NO: 117
Amino acid sequence of the 15H12/19D12 heavy chain B framework region 3	SEQ ID NO: 118
Nucleotide sequence encoding the 15H12/19D12 heavy chain B framework region 4	SEQ ID NO: 119
Amino acid sequence of the 15H12/19D12 heavy chain B framework region 4	SEQ ID NO: 120

CDR-L1 is the first complementarity determining region (CDR) which occurs in the light chain, CDR-L2 is the second CDR which occurs on the light chain and CDR-L3 is the third CDR which occurs on the light chain.

5 Similarly, CDR-H1 is the first CDR which occurs on the heavy chain, CDR-H2 is the second CDR which occurs on the heavy chain and CDR-H3 is the third CDR which occurs on the heavy chain.

FR-L1 is the first framework region of the light chain, FR-L2 is the second framework region of the light chain, FR-L3 is the third framework region of the light chain, FR-L4 is the fourth framework region on the light chain, FR-H1 is the first framework region of the heavy chain, FR-H2 is the second framework region of the heavy chain, FR-H3 is the third framework region of the heavy chain and FR-H4 is the fourth framework region of the heavy chain. These terms and the arrangement of CDRs and FRs on an immunoglobulin chain are well known in the art.

15 A mature light chain variable region of the invention, which lacks the signal peptide (*i.e.*, first 19 or 20 residues), is amino acids 20-128 of SEQ ID NO: 2, 41, 43, 72, 74, 76 or 78 which is encoded by nucleotides 58-384 of SEQ ID NO: 1, 40, 42, 71, 73, 75, or 77 or amino acids 21-130 of SEQ ID NO: 25 which is encoded by nucleotides 61-390 of SEQ ID NO: 24.

A mature heavy chain variable region, which lacks the signal peptide (*i.e.*, first 19 residues), is amino acids 20-137 of SEQ ID NO: 4, 45 or 112 which is encoded by nucleotides 58-411 of SEQ ID NO: 3, 44 or 111 or amino acids 20-140 of SEQ ID NO: 27 which is encoded by nucleotides 58-420 of SEQ ID NO: 26.

5 In some embodiments the 15H12 and 19D12 CDR-H1 is GFTFSSFAMH (SEQ ID NO: 17) which is encoded by the nucleotide sequence of SEQ ID NO: 18. In some embodiments the 1H3 CDR-H1 is NYAMH (SEQ ID NO: 70).

The present invention also includes antibodies and antigen-binding fragments which include the framework regions of the antibodies and antigen-binding fragments
10 of the invention. Preferably, FR-L1 is amino acids 20-42 of SEQ ID NO: 2 or amino acids 21-43 of SEQ ID NO: 25; FR-L2 is amino acids 54-68 of SEQ ID NO: 2 or amino acids 55-69 of SEQ ID NO: 25; FR-L3 is amino acids 76-107 of SEQ ID NO: 2 or amino acids 77-108 of SEQ ID NO: 25; FR-L4 is amino acids 117-128 of SEQ ID NO: 2 or amino acids 128-130 of SEQ ID NO: 25; FR-H1 is amino acids 20-44 or 20-
15 49 of SEQ ID NO: 4 or amino acids 20-44 or 20-49 of SEQ ID NO: 27; FR-H2 is amino acids 55-68 of SEQ ID NO: 4 or amino acids 55-68 of SEQ ID NO: 27; FR-H3 is amino acids 85-116 of SEQ ID NO: 4 or amino acids 85-116 of SEQ ID NO: 27 and FR-H4 is amino acids 127-137 of SEQ ID NO: 4 or amino acids 130-140 of SEQ ID NO: 27.

20 In preferred embodiments, the antibody molecules of the present invention include FR-L1 defined by amino acids 20-42 of SEQ ID NO: 41 or 43; FR-L2 defined by amino acids 54-68 of SEQ ID NO: 41 or 43; FR-L3 defined by amino acids 76-107 of SEQ ID NO: 41 or 43; and FR-L4 defined by amino acids 117-128 of SEQ ID NO: 41 or 43. Furthermore, preferred embodiments include antibody molecules including
25 FR-H1 defined by amino acids 20-44 of SEQ ID NO: 45; FR-H2 defined by amino acids 55-68 of SEQ ID NO: 45; FR-H3 defined by amino acids 85-116 of SEQ ID NO: 45; and FR-H4 defined by amino acids 127-137 of SEQ ID NO: 45.

Molecular Biology

30 In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, *e.g.*, Sambrook, Fritsch & Maniatis, Molecular Cloning: A Laboratory Manual, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (herein

"Sambrook, *et al.*, 1989"); DNA Cloning: A Practical Approach, Volumes I and II (D.N. Glover ed. 1985); Oligonucleotide Synthesis (M.J. Gait ed. 1984); Nucleic Acid Hybridization (B.D. Hames & S.J. Higgins eds. (1985)); Transcription And Translation (B.D. Hames & S.J. Higgins, eds. (1984)); Animal Cell Culture (R.I. Freshney, ed. (1986)); Immobilized Cells And Enzymes (IRL Press, (1986)); B. Perbal, A Practical Guide To Molecular Cloning (1984); F.M. Ausubel, *et al.* (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, Inc. (1994).

A "polynucleotide", "nucleic acid" or "nucleic acid molecule" may refer to the phosphate ester polymeric form of ribonucleosides (adenosine, guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine, deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules"), or any phosphoester analogs thereof, such as phosphorothioates and thioesters, in single stranded form, double-stranded form or otherwise.

A "polynucleotide sequence", "nucleic acid sequence" or "nucleotide sequence" is a series of nucleotide bases (also called "nucleotides") in a nucleic acid, such as DNA or RNA, and means any chain of two or more nucleotides.

A "coding sequence" or a sequence "encoding" an expression product, such as a RNA, polypeptide, protein, or enzyme, is a nucleotide sequence that, when expressed, results in production of the product.

The term "gene" means a DNA sequence that codes for or corresponds to a particular sequence of ribonucleotides or amino acids which comprise all or part of one or more RNA molecules, proteins or enzymes, and may or may not include regulatory DNA sequences, such as promoter sequences, which determine, for example, the conditions under which the gene is expressed. Genes may be transcribed from DNA to RNA which may or may not be translated into an amino acid sequence.

"Amplification" of DNA as used herein may denote the use of polymerase chain reaction (PCR) to increase the concentration of a particular DNA sequence within a mixture of DNA sequences. For a description of PCR see Saiki, *et al.*, Science (1988) 239: 487. In a specific embodiment, the present invention includes a nucleic acid, which encodes an anti-IGFR1 antibody, an anti-IGFR1 antibody heavy or light chain, an anti-IGFR1 antibody heavy or light chain variable region, an anti-IGFR1 antibody heavy or light chain constant region or anti-IGFR1 antibody CDR

(e.g., CDR- L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2 or CDR-H3) which can be amplified by PCR.

As used herein, the term "oligonucleotide" refers to a nucleic acid, generally of at least 10 (e.g., 10, 11, 12, 13 or 14), preferably at least 15 (e.g., 15, 16, 17, 18 or 19), and more preferably at least 20 nucleotides (e.g., 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30), preferably no more than 100 nucleotides (e.g., 40, 50, 60, 70, 80 or 90), that may be hybridizable to a genomic DNA molecule, a cDNA molecule, or an mRNA molecule encoding a gene, mRNA, cDNA, or other nucleic acid of interest.

Oligonucleotides can be labeled, e.g., by incorporation of ³²P-nucleotides, ³H-nucleotides, ¹⁴C-nucleotides, ³⁵S-nucleotides or nucleotides to which a label, such as biotin, has been covalently conjugated. In one embodiment, a labeled oligonucleotide can be used as a probe to detect the presence of a nucleic acid. In another embodiment, oligonucleotides (one or both of which may be labeled) can be used as PCR primers, either for cloning full length or a fragment of the gene, or to detect the presence of nucleic acids. Generally, oligonucleotides are prepared synthetically, preferably on a nucleic acid synthesizer.

The sequence of any nucleic acid (e.g., a nucleic acid encoding an IGFR1 gene or a nucleic acid encoding an anti-IGFR1 antibody or a fragment or portion thereof) may be sequenced by any method known in the art (e.g., chemical sequencing or enzymatic sequencing). "Chemical sequencing" of DNA may denote methods such as that of Maxam and Gilbert (1977) (Proc. Natl. Acad. Sci. USA 74:560), in which DNA is randomly cleaved using individual base-specific reactions. "Enzymatic sequencing" of DNA may denote methods such as that of Sanger (Sanger, *et al.*, (1977) Proc. Natl. Acad. Sci. USA 74:5463).

The nucleic acids herein may be flanked by natural regulatory (expression control) sequences, or may be associated with heterologous sequences, including promoters, internal ribosome entry sites (IRES) and other ribosome binding site sequences, enhancers, response elements, suppressors, signal sequences, polyadenylation sequences, introns, 5'- and 3'- non-coding regions, and the like.

A "promoter" or "promoter sequence" is a DNA regulatory region capable of binding an RNA polymerase in a cell (e.g., directly or through other promoter-bound proteins or substances) and initiating transcription of a coding sequence. A promoter sequence is, in general, bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or

elements necessary to initiate transcription at any level. Within the promoter sequence may be found a transcription initiation site (conveniently defined, for example, by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. The promoter may be operably associated with other expression control sequences, including enhancer and repressor sequences or with a nucleic acid of the invention (e.g., SEQ ID NO: 1, 3, 5-7, 11-13, 18, 22-24, 26, 28-30 or 34-36). Promoters which may be used to control gene expression include, but are not limited to, cytomegalovirus (CMV) promoter (U.S. Patent Nos. 5,385,839 and 5,168,062), the SV40 early promoter region (Benoist, *et al.*, (1981) *Nature* 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, *et al.*, (1980) *Cell* 22:787-797), the herpes thymidine kinase promoter (Wagner, *et al.*, (1981) *Proc. Natl. Acad. Sci. USA* 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster, *et al.*, (1982) *Nature* 296:39-42); prokaryotic expression vectors such as the β -lactamase promoter (Villa-Komaroff, *et al.*, (1978) *Proc. Natl. Acad. Sci. USA* 75:3727-3731), or the *tac* promoter (DeBoer, *et al.*, (1983) *Proc. Natl. Acad. Sci. USA* 80:21-25); see also "Useful proteins from recombinant bacteria" in *Scientific American* (1980) 242:74-94; and promoter elements from yeast or other fungi such as the *Gal 4* promoter, the *ADC* (alcohol dehydrogenase) promoter, *PGK* (phosphoglycerol kinase) promoter or the alkaline phosphatase promoter.

A coding sequence is "under the control of", "functionally associated with" or "operably associated with" transcriptional and translational control sequences in a cell when the sequences direct RNA polymerase mediated transcription of the coding sequence into RNA, preferably mRNA, which then may be trans-RNA spliced (if it contains introns) and, optionally, translated into a protein encoded by the coding sequence.

The terms "express" and "expression" mean allowing or causing the information in a gene, RNA or DNA sequence to become manifest; for example, producing a protein by activating the cellular functions involved in transcription and translation of a corresponding gene. A DNA sequence is expressed in or by a cell to form an "expression product" such as an RNA (e.g., mRNA) or a protein (e.g., antibody 1H3, 15H12 or 19D12 or a fragment thereof). The expression product itself may also be said to be "expressed" by the cell.

The terms "vector", "cloning vector" and "expression vector" mean the vehicle (e.g., a plasmid) by which a DNA or RNA sequence can be introduced into a host cell, so as to transform the host and, optionally, promote expression and/or replication of the introduced sequence.

5 The term "transfection" or "transformation" means the introduction of a nucleic acid into a cell. These terms may refer to the introduction of a nucleic acid encoding an anti-IGFR1 antibody or fragment thereof into a cell. The introduced gene or sequence may be called a "clone". A host cell that receives the introduced DNA or RNA has been "transformed" and is a "transformant" or a "clone". The DNA or RNA
10 introduced to a host cell can come from any source, including cells of the same genus or species as the host cell, or cells of a different genus or species.

 The term "host cell" means any cell of any organism that is selected, modified, transfected, transformed, grown, or used or manipulated in any way, for the production of a substance by the cell, for example the expression or replication, by
15 the cell, of a gene, a DNA or RNA sequence, a protein or an enzyme.

 The term "expression system" means a host cell and compatible vector which, under suitable conditions, can express a protein or nucleic acid which is carried by the vector and introduced to the host cell. Common expression systems include *E. coli* host cells and plasmid vectors, insect host cells and Baculovirus vectors, and
20 mammalian host cells and vectors. In a specific embodiment, IGFR1 or an antibody and antigen-binding fragment of the invention may be expressed in human embryonic kidney cells (HEK293). Other suitable cells include CHO (chinese hamster ovary) cells, HeLa cells and NIH 3T3 cells and NSO cells (non-Ig-producing murine myeloma cell line). Nucleic acids encoding an antibody or antigen-binding fragment of the
25 invention, sIGFR1 or IGFR1 may be expressed at high levels in an *E.coli*/T7 expression system as disclosed in U.S. Patent Nos. 4,952,496, 5,693,489 and 5,869,320 and in Davanloo, P., *et al.*, (1984) Proc. Natl. Acad. Sci. USA 81, 2035-2039; Studier, F. W., *et al.*, (1986) J. Mol. Biol. 189: 113-130; Rosenberg, A. H., *et al.*, (1987) Gene 56: 125-135; and Dunn, J. J., *et al.*, (1988) Gene 68: 259 which are
30 herein incorporated by reference.

 The present invention contemplates any superficial or slight modification to the amino acid or nucleotide sequences which correspond to the antibodies or antigen-binding fragments of the invention. In particular, the present invention contemplates sequence conservative variants of the nucleic acids which encode the antibodies or

antigen-binding fragments of the invention. "Sequence-conservative variants" of a polynucleotide sequence are those in which a change of one or more nucleotides in a given codon results in no alteration in the amino acid encoded at that position.

Function-conservative variants of the antibodies of the invention are also

- 5 contemplated by the present invention. "Function-conservative variants" are those in which one or more amino acid residues in a protein or enzyme have been changed without altering the overall conformation and function of the polypeptide, including, but, by no means, limited to, replacement of an amino acid with one having similar properties. Amino acids with similar properties are well known in the art. For
- 10 example, polar/hydrophilic amino acids which may be interchangeable include asparagine, glutamine, serine, cysteine, threonine, lysine, arginine, histidine, aspartic acid and glutamic acid; nonpolar/hydrophobic amino acids which may be interchangeable include glycine, alanine, valine, leucine, isoleucine, proline, tyrosine, phenylalanine, tryptophan and methionine; acidic amino acids which may be
- 15 interchangeable include aspartic acid and glutamic acid and basic amino acids which may be interchangeable include histidine, lysine and arginine.

- The present invention includes anti-IGFR1 antibodies and fragments thereof which are encoded by nucleic acids as described in Table 1 as well as nucleic acids which hybridize thereto. Preferably, the nucleic acids hybridize under low stringency
- 20 conditions, more preferably under moderate stringency conditions and most preferably under high stringency conditions and, preferably, exhibit IGFR1 binding activity. A nucleic acid molecule is "hybridizable" to another nucleic acid molecule, such as a cDNA, genomic DNA, or RNA, when a single stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate
- 25 conditions of temperature and solution ionic strength (see Sambrook, *et al.*, *supra*). The conditions of temperature and ionic strength determine the "stringency" of the hybridization. Typical low stringency hybridization conditions may be 55°C, 5X SSC, 0.1% SDS, 0.25% milk, and no formamide; or 30% formamide, 5X SSC, 0.5% SDS. Typical, moderate stringency hybridization conditions are similar to the low stringency
- 30 conditions except the hybridization is carried out in 40% formamide, with 5X or 6X SSC. High stringency hybridization conditions are similar to low stringency conditions except the hybridization conditions are carried out in 50% formamide, 5X or 6X SSC and, optionally, at a higher temperature (*e.g.*, 57°C, 59°C, 60°C, 62°C, 63°C, 65°C or 68°C). In general, SSC is 0.15M NaCl and 0.015M Na-citrate. Hybridization

requires that the two nucleic acids contain complementary sequences, although, depending on the stringency of the hybridization, mismatches between bases are possible. The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater the degree of similarity or homology between two nucleotide sequences, the higher the stringency under which the nucleic acids may hybridize. For hybrids of greater than 100 nucleotides in length, equations for calculating the melting temperature have been derived (see Sambrook, *et al.*, *supra*, 9.50-9.51). For hybridization with shorter nucleic acids, *i.e.*, oligonucleotides, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook, *et al.*, *supra*, 11.7-11.8).

Also included in the present invention are nucleic acids comprising nucleotide sequences and polypeptides comprising amino acid sequences which are at least about 70% identical, preferably at least about 80% identical, more preferably at least about 90% identical and most preferably at least about 95% identical (*e.g.*, 95%, 96%, 97%, 98%, 99%, 100%) to the reference nucleotide and amino acid sequences of Table 1 when the comparison is performed by a BLAST algorithm wherein the parameters of the algorithm are selected to give the largest match between the respective sequences over the entire length of the respective reference sequences. Polypeptides comprising amino acid sequences which are at least about 70% similar, preferably at least about 80% similar, more preferably at least about 90% similar and most preferably at least about 95% similar (*e.g.*, 95%, 96%, 97%, 98%, 99%, 100%) to the reference amino acid sequences of Table 1 (*e.g.*, SEQ ID NOs. 2 (*e.g.*, amino acids 20-128), 4 (*e.g.*, amino acids 20-137), 8-10, 14-16, 17, 25 (*e.g.*, amino acids 21-130), 27 (*e.g.*, amino acids 20-140), 31-33 or 37-39) when the comparison is performed with a BLAST algorithm wherein the parameters of the algorithm are selected to give the largest match between the respective sequences over the entire length of the respective reference sequences, are also included in the present invention.

Sequence identity refers to exact matches between the nucleotides or amino acids of two sequences which are being compared. Sequence similarity refers to both exact matches between the amino acids of two polypeptides which are being compared in addition to matches between nonidentical, biochemically related amino

acids. Biochemically related amino acids which share similar properties and may be interchangeable are discussed above.

- The following references regarding the BLAST algorithm are herein incorporated by reference: **BLAST ALGORITHMS**: Altschul, S.F., *et al.*, (1990) J. Mol. Biol. 215:403-410; Gish, W., *et al.*, (1993) Nature Genet. 3:266-272; Madden, T.L., *et al.*, (1996) Meth. Enzymol. 266:131-141; Altschul, S.F., *et al.*, (1997) Nucleic Acids Res. 25:3389-3402; Zhang, J., *et al.*, (1997) Genome Res. 7:649-656; Wootton, J.C., *et al.*, (1993) Comput. Chem. 17:149-163; Hancock, J.M. *et al.*, (1994) Comput. Appl. Biosci. 10:67-70; **ALIGNMENT SCORING SYSTEMS**: Dayhoff, M.O., *et al.*, "A model of evolutionary change in proteins." in Atlas of Protein Sequence and Structure, (1978) vol. 5, suppl. 3. M.O. Dayhoff (ed.), pp. 345-352, Natl. Biomed. Res. Found., Washington, DC; Schwartz, R.M., *et al.*, "Matrices for detecting distant relationships." in Atlas of Protein Sequence and Structure, (1978) vol. 5, suppl. 3. M.O. Dayhoff (ed.), pp. 353-358, Natl. Biomed. Res. Found., Washington, DC; Altschul, S.F., (1991) J. Mol. Biol. 219:555-565; States, D.J., *et al.*, (1991) Methods 3:66-70; Henikoff, S., *et al.*, (1992) Proc. Natl. Acad. Sci. USA 89:10915-10919; Altschul, S.F., *et al.*, (1993) J. Mol. Evol. 36:290-300; **ALIGNMENT STATISTICS**: Karlin, S., *et al.*, (1990) Proc. Natl. Acad. Sci. USA 87:2264-2268; Karlin, S., *et al.*, (1993) Proc. Natl. Acad. Sci. USA 90:5873-5877; Dembo, A., *et al.*, (1994) Ann. Prob. 22:2022-2039; and Altschul, S.F. "Evaluating the statistical significance of multiple distinct local alignments." in Theoretical and Computational Methods in Genome Research (S. Suhai, ed.), (1997) pp. 1-14, Plenum, New York.

Antibody Structure

- In general, the basic antibody structural unit is known to comprise a tetramer. Each tetramer includes two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain may include a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain may define a constant region primarily responsible for effector function. Typically, human light chains are classified as kappa and lambda light chains. Furthermore, human heavy chains are typically classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. Within light and heavy chains, the variable and constant

regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids. See generally, Fundamental Immunology Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)) (incorporated by reference in its entirety for all purposes).

5 The variable regions of each light/heavy chain pair may form the antibody binding site. Thus, in general, an intact IgG antibody has two binding sites. Except in bifunctional or bispecific antibodies, the two binding sites are, in general, the same.

Normally, the chains all exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called
10 complementarity determining regions or CDRs. The CDRs from the two chains of each pair are usually aligned by the framework regions, enabling binding to a specific epitope. In general, from N-terminal to C-terminal, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is, generally, in accordance with the
15 definitions of Sequences of Proteins of Immunological Interest, Kabat, *et al.*; National Institutes of Health, Bethesda, Md. ; 5th ed.; NIH Publ. No. 91-3242 (1991); Kabat (1978) Adv. Prot. Chem. 32:1-75; Kabat, *et al.*, (1977) J. Biol. Chem. 252:6609-6616; Chothia, *et al.*, (1987) J Mol. Biol. 196:901-917 or Chothia, *et al.*, (1989) Nature 342:878-883. The present invention provides antibodies or antigen-binding
20 fragments of the invention comprising CDRs and FRs from the light and heavy chains of 1H3, 15H12 and 19D12 (*e.g.*, 15H12/19D12 LCA, 15H12/19D12 LCB, 15H12/19D12 HCA, SEQ ID NOs: 2, 4, 25, 27, 41, 43 and 45) as defined by Kabat and Chothia (see above references).

25 **Antibody Molecules**

The term "antibody molecule" includes, but is not limited to, antibodies and fragments, preferably antigen-binding fragments, thereof. The term includes monoclonal antibodies, polyclonal antibodies, bispecific antibodies, Fab antibody fragments, F(ab)₂ antibody fragments, Fv antibody fragments (*e.g.*, V_H or V_L), single
30 chain Fv antibody fragments and dsFv antibody fragments. Furthermore, the antibody molecules of the invention may be fully human antibodies or chimeric antibodies. Preferably, the antibody molecules are monoclonal, fully human antibodies; more preferably, the antibody molecules are 1H3, 15H12 or 19D12.

Preferably, the antibody molecules include one or more of the variable regions and CDRs whose amino acid and nucleotide sequences are set forth in Table 1.

The present invention includes any antibody molecule comprising a CDR selected from:

- 5 RASQSIGSSLH (SEQ ID NO: 8);
YASQSL (SEQ ID NO: 9);
HQSSRLPHT (SEQ ID NO: 10);
SFAMH (SEQ ID NO: 14)
GFTFSSFAMH (SEQ ID NO: 17);
- 10 VIDTRGATYYADSVKG (SEQ ID NO: 15);
LGNFYYGMDV (SEQ ID NO: 16);
RASQSVSSFLA (SEQ ID NO: 31);
DASNRAP (SEQ ID NO: 32);
QQRSNWPRWT (SEQ ID NO: 33);
- 15 GFTFSNYAMH (SEQ ID NO: 37);
AIGAGGDTYYADSVKG (SEQ ID NO: 38); and
GRHRNWYYYNKDY (SEQ ID NO: 39);
NYAMH (SEQ ID NO: 70)

The scope of the present invention includes antibody variable regions of the present invention (*e.g.*, any variable region, mature or unprocessed, indicated in Table 1) linked to any immunoglobulin constant region. If a light chain variable region is linked to a constant region, preferably it is a κ chain. If a heavy chain variable region is linked to a constant region, preferably it is a γ 1, γ 2, γ 3 or γ 4 constant region, more preferably, γ 1, γ 2 or γ 4 and even more preferably γ 1 or γ 4.

The anti-IGFR1 antibody molecules of the invention preferably recognize human IGFR1, preferably sIGFR1; however, the present invention includes antibody molecules which recognize IGFR1 from different species, preferably mammals (*e.g.*, mouse, rat, rabbit, sheep or dog). The present invention also includes anti-IGFR1 antibodies or fragments thereof which are complexed with IGFR1 or any fragment thereof (*e.g.*, amino acids 30-902 of SEQ ID NO: 19) or with any cell which is expressing IGFR1 or any portion or fragment thereof on the cell surface (*e.g.*, HEK293 cells stably transformed with human *IGFR1* or MCF7 (*e.g.*, ATCC Cell Line No. HTB-22)). Such complexes may be made by contacting the antibody or antibody fragment with IGFR1 or the IGFR1 fragment.

In a preferred embodiment, fully-human monoclonal antibodies directed against IGFR1 are generated using transgenic mice carrying parts of the human immune system rather than the mouse system. These transgenic mice, which may be referred to, herein, as "HuMAb" mice, contain a human immunoglobulin gene miniloci that encodes unrearranged human heavy (μ and γ) and κ light chain immunoglobulin sequences, together with targeted mutations that inactivate the

endogenous μ and κ chain loci (Lonberg, N., *et al.*, (1994) *Nature* 368(6474): 856-859). Accordingly, the mice exhibit reduced expression of mouse IgM or κ , and in response to immunization, the introduced human heavy and light chain transgenes undergo class switching and somatic mutation to generate high affinity human IgG κ monoclonal antibodies (Lonberg, N., *et al.*, (1994), *supra*; reviewed in Lonberg, N. (1994) *Handbook of Experimental Pharmacology* 113:49-101; Lonberg, N., *et al.*, (1995) *Intern. Rev. Immunol.* 13:65-93, and Harding, F., *et al.*, (1995) *Ann. N. Y Acad. Sci* 764:536-546). The preparation of HuMab mice is commonly known in the art and is described, for example, in Taylor, L., *et al.*, (1992) *Nucleic Acids Research* 20:6287-6295; Chen, J., *et al.*, (1993) *International Immunology* 5: 647-656; Tuaillon, *et al.*, (1993) *Proc. Natl. Acad. Sci USA* 90:3720-3724; Choi, *et al.*, (1993) *Nature Genetics* 4:117-123; Chen, J., *et al.*, (1993) *EMBO J.* 12: 821- 830; Tuaillon, *et al.*, (1994) *J Immunol.* 152:2912-2920; Lonberg, *et al.*, (1994) *Nature* 368(6474): 856-859; Lonberg, N. (1994) *Handbook of Experimental Pharmacology* 113:49-101; Taylor, L., *et al.*, (1994) *International Immunology* 6: 579-591; Lonberg, N., *et al.*, (1995) *Intern. Rev. Immunol. Vol. 13: 65-93*; Harding, F., *et al.*, (1995) *Ann. N.Y Acad. Sci* 764:536-546; Fishwild, D., *et al.*, (1996) *Nature Biotechnology* 14: 845-851 and Harding, *et al.*, (1995) *Annals NY Acad. Sci.* 764:536-546; the contents of all of which are hereby incorporated by reference in their entirety. See further, U.S. Patent Nos. 5,545,806; 5, 569,825; 5,625,126; 5,633,425; 5,789,650; 5,877,397; 5,661,016; 5,814,318; 5,874, 299; 5,770,429 and 5,545,807; and International Patent Application Publication Nos. WO 98/24884; WO 94/25585; WO 93/12227; WO 92/22645 and WO 92/03918 the disclosures of all of which are hereby incorporated by reference in their entity.

To generate fully human, monoclonal antibodies to IGFR1, HuMab mice can be immunized with an antigenic IGFR1 polypeptide, preferably amino acids 30-902 of SEQ ID NO: 19, as described by Lonberg, N., *et al.*, (1994) *Nature* 368(6474): 856-859; Fishwild, D., *et al.*, (1996) *Nature Biotechnology* 14: 845-851 and WO 98/24884. Preferably, the mice will be 6-16 weeks of age upon the first immunization. For example, a purified preparation of IGFR1 or sIGFR1 can be used to immunize the HuMab mice intraperitoneally. The mice can also be immunized with whole HEK293 cells which are stably transformed or transfected with an *IGFR1* gene. An "antigenic IGFR1 polypeptide" may refer to an IGFR1 polypeptide of any fragment thereof,

preferably amino acids 30-902 of SEQ ID NO: 19, which elicits an anti-IGFR1 immune response, preferably in HuMAb mice.

In general, HuMAb transgenic mice respond well when initially immunized intraperitoneally (IP) with antigen in complete Freund's adjuvant, followed by every
5 other week IP immunizations (usually, up to a total of 6) with antigen in incomplete Freund's adjuvant. Mice can be immunized, first, with cells expressing IGFR1 (e.g., stably transformed HEK293 cells), then with a soluble fragment of IGFR1 (e.g., amino acids 30-902 of SEQ ID NO: 19) and continually receive alternating immunizations with the two antigens. The immune response can be monitored over the course of
10 the immunization protocol with plasma samples being obtained by retroorbital bleeds. The plasma can be screened for the presence of anti-IGFR1 antibodies, for example by ELISA, and mice with sufficient titers of immunoglobulin can be used for fusions. Mice can be boosted intravenously with antigen 3 days before sacrifice and removal of the spleen. It is expected that 2-3 fusions for each antigen may need to be
15 performed. Several mice can be immunized for each antigen. For example, a total of twelve HuMAb mice of the HC07 and HC012 strains can be immunized.

Hybridoma cells which produce the monoclonal, fully human anti-IGFR1 antibodies may be produced by methods which are commonly known in the art. These methods include, but are not limited to, the hybridoma technique originally
20 developed by Kohler, *et al.*, (1975) (Nature 256:495-497), as well as the trioma technique (Hering, *et al.*, (1988) Biomed. Biochim. Acta. 47:211-216 and Hagiwara, *et al.*, (1993) Hum. Antibod. Hybridomas 4:15), the human B-cell hybridoma technique (Kozbor, *et al.*, (1983) Immunology Today 4:72 and Cote, *et al.*, (1983) Proc. Natl. Acad. Sci. U.S.A 80:2026-2030), and the EBV-hybridoma technique (Cole, *et al.*, in
25 Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96, 1985). Preferably, mouse splenocytes are isolated and fused with PEG to a mouse myeloma cell line based upon standard protocols. The resulting hybridomas may then be screened for the production of antigen-specific antibodies. For example, single cell suspensions of splenic lymphocytes from immunized mice may be fused to one-sixth
30 the number of P3X63- Ag8.653 nonsecreting mouse myeloma cells (ATCC, CRL 1580) with 50% PEG. Cells may be plated at approximately 2×10^5 cells/mL in a flat bottom microtiter plate, followed by a two week incubation in selective medium containing 20% fetal Clone Serum, 18% "653" conditioned media, 5% origen (IGEN), 4 mM L-glutamine, 1 mM L-glutamine, 1 mM sodium pyruvate, 5mM HEPES, 0.055

mM 2-mercaptoethanol, 50 units/ml penicillin, 50 mg/ml streptomycin, 50 mg/ml gentamycin and 1X HAT (Sigma; the HAT is added 24 hours after the fusion). After two weeks, cells may be cultured in medium in which the HAT is replaced with HT. Individual wells may then be screened by ELISA for human anti-IGFR1 monoclonal
5 IgG antibodies. Once extensive hybridoma growth occurs, medium can be observed usually after 10-14 days. The antibody secreting hybridomas may be replated, screened again, and if still positive for human IgG, anti-IGFR1 monoclonal antibodies, can be subcloned at least twice by limiting dilution. The stable subclones may then be cultured *in vitro* to generate small amounts of antibody in tissue culture medium for
10 characterization.

The anti-IGFR antibody molecules of the present invention may also be produced recombinantly (*e.g.*, in an *E.coli*/T7 expression system as discussed above). In this embodiment, nucleic acids encoding the antibody molecules of the invention (*e.g.*, V_H or V_L) may be inserted into a pET-based plasmid and expressed in
15 the *E.coli*/T7 system. There are several methods by which to produce recombinant antibodies which are known in the art. One example of a method for recombinant production of antibodies is disclosed in U.S. Patent No. 4,816,567 which is herein incorporated by reference. Transformation can be by any known method for
introducing polynucleotides into a host cell. Methods for introduction of heterologous
20 polynucleotides into mammalian cells are well known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene-mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, biolistic injection and direct microinjection of the DNA into nuclei. In addition, nucleic acid molecules may be introduced into mammalian cells by viral
25 vectors. Methods of transforming cells are well known in the art. See, for example, U.S. Patent Nos. 4,399,216; 4,912,040; 4,740,461 and 4,959,455.

Mammalian cell lines available as hosts for expression are well known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC). These include, *inter alia*, Chinese hamster ovary (CHO)
30 cells, NSO, SP2 cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (*e.g.*, Hep G2), A549 cells, 3T3 cells, and a number of other cell lines. Mammalian host cells include human, mouse, rat, dog, monkey, pig, goat, bovine, horse and hamster cells. Cell lines of particular preference are selected through determining which cell lines have high expression

levels. Other cell lines that may be used are insect cell lines, such as Sf9 cells, amphibian cells, bacterial cells, plant cells and fungal cells. When recombinant expression vectors encoding the heavy chain or antigen-binding portion thereof, the light chain and/or antigen-binding portion thereof are introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably, secretion of the antibody into the culture medium in which the host cells are grown.

Antibodies can be recovered from the culture medium using standard protein purification methods. Further, expression of antibodies of the invention (or other moieties therefrom) from production cell lines can be enhanced using a number of known techniques. For example, the glutamine synthetase gene expression system (the GS system) is a common approach for enhancing expression under certain conditions. The GS system is discussed in whole or part in connection with European Patent Nos. 0 216 846, 0 256 055, and 0 323 997 and European Patent Application No. 89303964.4.

It is likely that antibodies expressed by different cell lines or in transgenic animals will have different glycosylation from each other. However, all antibodies encoded by the nucleic acid molecules provided herein, or comprising the amino acid sequences provided herein are part of the instant invention, regardless of the glycosylation of the antibodies.

" K_{off} " refers to the off-rate constant for dissociation of the antibody from an antibody/antigen complex.

" K_{on} " refers to the rate at which the antibody associates with the antigen.

" K_d " refers to the dissociation constant of a particular antibody/antigen interaction. $K_d = K_{off}/K_{on}$.

The term "monoclonal antibody," as used herein, refers to an antibody obtained from a population of substantially homogeneous antibodies, *i.e.*, the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Monoclonal antibodies are advantageous in that they may be synthesized by a hybridoma culture, essentially uncontaminated by other immunoglobulins. The modifier "monoclonal" indicates the character of the antibody as being amongst a substantially homogeneous population of antibodies, and is not to be construed as

requiring production of the antibody by any particular method. As mentioned above, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler, *et al.*, (1975) Nature 256: 495.

5 A polyclonal antibody is an antibody which was produced among or in the presence of one or more other, non-identical antibodies. In general, polyclonal antibodies are produced from a B-lymphocyte in the presence of several other B-lymphocytes which produced non-identical antibodies. Usually, polyclonal antibodies are obtained directly from an immunized animal.

10 A bispecific or bifunctional antibody is an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Bispecific antibodies can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, *e.g.*, Songsivilai, *et al.*, (1990) Clin. Exp. Immunol. 79: 315-321, Kostelny, *et al.*, (1992) J Immunol. 148:1547- 1553. In addition, bispecific antibodies
15 may be formed as "diabodies" (Holliger, *et al.*, (1993) PNAS USA 90:6444-6448) or as "Janusins" (Traunecker, *et al.*, (1991) EMBO J. 10:3655-3659 and Traunecker, *et al.*, (1992) Int. J. Cancer Suppl. 7:51-52).

 The term "fully human antibody" refers to an antibody which comprises human immunoglobulin protein sequences only. A fully human antibody may contain murine
20 carbohydrate chains if produced in a mouse, in a mouse cell or in a hybridoma derived from a mouse cell. Similarly, "mouse antibody" refers to an antibody which comprises mouse immunoglobulin sequences only.

 The present invention includes "chimeric antibodies"- an antibody which comprises a variable region of the present invention fused or chimerized with an
25 antibody region (*e.g.*, constant region) from another, non-human species (*e.g.*, mouse, horse, rabbit, dog, cow, chicken). These antibodies may be used to modulate the expression or activity of IGFR1 in the non-human species.

 "Single-chain Fv" or "sFv" antibody fragments have the V_H and V_L domains of an antibody, wherein these domains are present in a single polypeptide chain.
30 Generally, the sFv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. Techniques described for the production of single chain antibodies (U.S. Patent Nos. 5,476,786; 5,132,405 and 4,946,778) can be adapted to produce anti-IGFR1-specific single chain antibodies. For a review of sFv see Pluckthun in The

Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds. Springer-Verlag, N.Y., pp. 269-315 (1994).

“Disulfide stabilized Fv fragments” and “dsFv” refer to antibody molecules comprising a variable heavy chain (V_H) and a variable light chain (V_L) which are linked by a disulfide bridge.

Antibody fragments within the scope of the present invention also include $F(ab)_2$ fragments which may be produced by enzymatic cleavage of an IgG by, for example, pepsin. Fab fragments may be produced by, for example, reduction of $F(ab)_2$ with dithiothreitol or mercaptoethylamine. A Fab fragment is a V_L - C_L chain appended to a V_H - C_{H1} chain by a disulfide bridge. A $F(ab)_2$ fragment is two Fab fragments which, in turn, are appended by two disulfide bridges. The Fab portion of an $F(ab)_2$ molecule includes a portion of the F_c region between which disulfide bridges are located.

An F_v fragment is a V_L or V_H region.

Depending on the amino acid sequences of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are at least five major classes of immunoglobulins: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), *e.g.* IgG-1, IgG-2, IgG-3 and IgG-4; IgA-1 and IgA-2.

The anti-IGFR1 antibody molecules of the invention may also be conjugated to a chemical moiety. The chemical moiety may be, *inter alia*, a polymer, a radionuclide or a cytotoxic factor. Preferably the chemical moiety is a polymer which increases the half-life of the antibody molecule in the body of a subject. Suitable polymers include, but are not limited to, polyethylene glycol (PEG) (*e.g.*, PEG with a molecular weight of 2kDa, 5 kDa, 10 kDa, 12kDa, 20 kDa, 30kDa or 40kDa), dextran and monomethoxypolyethylene glycol (mPEG). Lee, *et al.*, (1999) (Bioconj. Chem. 10:973-981) discloses PEG conjugated single-chain antibodies. Wen, *et al.*, (2001) (Bioconj. Chem. 12:545-553) disclose conjugating antibodies with PEG which is attached to a radiometal chelator (diethylenetriaminopentaacetic acid (DTPA)).

The antibodies and antibody fragments of the invention may also be conjugated with labels such as ^{99}Tc , ^{90}Y , ^{111}In , ^{32}P , ^{14}C , ^{125}I , ^3H , ^{131}I , ^{11}C , ^{15}O , ^{13}N , ^{18}F , ^{35}S , ^{51}Cr , ^{57}To , ^{226}Ra , ^{60}Co , ^{59}Fe , ^{57}Se , ^{152}Eu , ^{67}Cu , ^{217}Ci , ^{211}At , ^{212}Pb , ^{47}Sc , ^{109}Pd , ^{234}Th , and ^{40}K , ^{157}Gd , ^{55}Mn , ^{52}Tr and ^{56}Fe .

The antibodies and antibody fragments of the invention may also be conjugated with fluorescent or chemiluminescent labels, including fluorophores such as rare earth chelates, fluorescein and its derivatives, rhodamine and its derivatives, isothiocyanate, phycoerythrin, phycocyanin, allophycocyanin, o-phthalaldehyde, fluorescamine, ¹⁵²Eu, dansyl, umbelliferone, luciferin, luminal label, isoluminal label, an aromatic acridinium ester label, an imidazole label, an acridinium salt label, an oxalate ester label, an aequorin label, 2,3-dihydrophthalazinediones, biotin/avidin, spin labels and stable free radicals.

The antibody molecules may also be conjugated to a cytotoxic factor such as diphtheria toxin, *Pseudomonas aeruginosa* exotoxin A chain, ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins and compounds (e.g., fatty acids), dianthin proteins, *Phytolacca americana* proteins PAPI, PAPII, and PAP-S, *momordica charantia* inhibitor, curcin, crotin, *saponaria officinalis* inhibitor, mitogellin, restrictocin, phenomycin, and enomycin.

Any method known in the art for conjugating the antibody molecules of the invention to the various moieties may be employed, including those methods described by Hunter, *et al.*, (1962) Nature 144:945; David, *et al.*, (1974) Biochemistry 13:1014; Pain, *et al.*, (1981) J. Immunol. Meth. 40:219; and Nygren, J., (1982) Histochem. and Cytochem. 30:407. Methods for conjugating antibodies are conventional and very well known in the art.

Modified Antibody Molecules

The present invention includes antibodies and antigen-binding fragments (e.g., fully-human antibodies, SFv, dsFv, Fv, chimeric antibodies) comprising a light chain of SEQ ID NOs: 41, 43, 72, 74, 76 or 78 (15H12/19D12 LCA, LCB, LCC, LCD, LCE or LCF); preferably amino acids 20-128 of SEQ ID NOs: 41, 43, 72, 74, 76 or 78 (mature 15H12/19D12 LCA, LCB, LCC, LCD, LCE or LCF). The present invention also includes antibody molecules including the heavy chain of SEQ ID NO: 45 or 112 (15H12/19D12 HCA, HCB); preferably amino acids 20-137 of SEQ ID NO: 45 or 112 (mature 15H12/19D12 HCA, HCB).

The 15H12/19D12 LCA, LCB, LCC, LCD, LCE and LCF may be dimerized with any other immunoglobulin heavy chain, preferably an immunoglobulin heavy chain of the present invention. Likewise, 15H12/19D12 HCA or HCB may be dimerized with any light chain, preferably a light chain of the present invention. For example,

15H12/19D12 HCA or HCB may be dimerized with 15H12/19D12 LCC, LCD, LCE or LCF.

Antibodies and antigen-binding fragments comprising 15H12/19D12 LCA,
15H12/19D12 LCB, 15H12/19D12 LCC, 15H12/19D12 LCD, 15H12/19D12 LCE,
5 15H12/19D12 LCF, 15H12/19D12 HCA or 15H12/19D12 HCB or any fragment
thereof exhibit minimal immunogenicity in a human subject; thereby, leading to a low
incidence of HAHA response when administered to a human subject.

Preferred antibody chains are shown below. Dotted underscored type
encodes the signal peptide. Solid underscored type encodes the CDRs. Plain type
10 encodes the framework regions. Most preferably, the antibody chains are mature
fragments which lack the signal peptide.

Modified 19D12/15H12 Light Chain-C (SEQ ID NO: 71)

15 ATG TCG CCA TCA CAA CTC ATT GGG TTT CTG CTG CTC TGG GTT CCA GCC TCC
 AGG GGT GAA ATT GTG CTG ACT CAG AGC CCA GAC TCT CTG TCT GTG ACT CCA
 GGC GAG AGA GTC ACC ATC ACC TGC CCG GCC AGT CAG AGC ATT GGT AGT AGC
20 TTA CAC TGG TAC CAG CAG AAA CCA GGT CAG TCT CCA AAG CTT CTC ATC AAG
 TAT GCA TCC CAG TCC CTC TCA GGG GTC CCC TCG AGG TTC AGT GGC AGT GGA
25 TCT GGG ACA GAT TTC ACC CTC ACC ATC AGT AGC CTC GAG GCT GAA GAT GCT
 GCA GCG TAT TAC TGT CAT CAG AGT AGT CGT TTA CCT CAC ACT TTC GGC CAA
 GGG ACC AAG GTG GAG ATC AAA CGT ACG

30

(SEQ ID NO: 72)

 M S P S Q L I G F L L L W V P A S
35 R G E I V L T Q S P D S L S V T P
 G E R V T I T C R A S Q S I G S S
 L H W Y Q Q K P G Q S P K L L I K
40 Y A S Q S L S G V P S R F S G S G
 S G T D F T L T I S S L E A E D A
45 A A Y Y C H Q S S R L P H T F G Q
 G T K V E I K R T

Modified 19D12/15H12 Light Chain-D (SEQ ID NO: 73)

5 ATG TCG CCA TCA CAA CTC ATT GGG TTT CTG CTG CTC TGG GTT CCA GCC TCC
 AGG GGT GAA ATT GTG CTG ACT CAG AGC CCA GAC TCT CTG TCT GTG ACT CCA
 GGC GAG AGA GTC ACC ATC ACC TGC CGG GCC AGT CAG AGC ATT GGT AGT AGC
 10 TTA CAC TGG TAC CAG CAG AAA CCA GGT CAG TCT CCA AAG CTT CTC ATC AAG
 TAT GCA TCC CAG TCC CTC TCA GGG GTC CCC TCG AGG TTC AGT GGC AGT GGA
 TCT GGG ACA GAT TTC ACC CTC ACC ATC AGT AGC CTC GAG GCT GAA GAT TTC
 15 GCA GTG TAT TAC TGT CAT CAG AGT AGT CGT TTA CCT CAC ACT TTC GGC CAA
 GGG ACC AAG GTG GAG ATC AAA CGT ACG

(SEQ ID NO: 74)

20 M S P S Q L I G F L L L W V P A S
 R G E I V L T Q S P D S L S V T P
 25 G E R V T I T C R A S Q S I G S S
 L H W Y Q Q K P G Q S P K L L I K
 30 Y A S Q S L S G V P S R F S G S G
 S G T D F T L T I S S L E A E D F
 A V Y Y C H Q S S R L P H T F G Q
 35 G T K V E I K R T

Modified 19D12/15H12 Light Chain-E (SEQ ID NO: 75)

40 ATG TCG CCA TCA CAA CTC ATT GGG TTT CTG CTG CTC TGG GTT CCA GCC TCC
 AGG GGT GAA ATT GTG CTG ACT CAG AGC CCA GGT ACC CTG TCT GTG TCT CCA
 GGC GAG AGA GCC ACC CTC TCC TGC CGG GCC AGT CAG AGC ATT GGT AGT AGC
 45 TTA CAC TGG TAC CAG CAG AAA CCA GGT CAG GCT CCA AGG CTT CTC ATC AAG
 TAT GCA TCC CAG TCC CTC TCA GGG ATC CCC GAT AGG TTC AGT GGC AGT GGA
 TCT GGG ACA GAT TTC ACC CTC ACC ATC AGT AGA CTG GAG CCT GAA GAT GCT
 50 GCA GCG TAT TAC TGT CAT CAG AGT AGT CGT TTA CCT CAC ACT TTC GGC CAA
 GGG ACC AAG GTG GAG ATC AAA CGT ACA

55 (SEQ ID NO: 76)

5 M S P S Q L I G F L L L W V P A S
 R G E I V L T Q S P G T L S V S P
 G E R A T L S C R A S Q S I G S S
 10 L H W Y Q Q K P G Q A P R L L I K
 Y A S Q S L S G I P D R F S G S G
 S G T D F T L T I S R L E P E D A
 15 A A Y Y C H Q S S R L P H T F G Q
 G T K V E I K R T

Modified 19D12/15H12 Light Chain-F (SEQ ID NO: 77)

20 ATG TCG CCA TCA CAA CTC ATT GGG TTT CTG CTG CTC TGG GTT CCA GCC TCC
 AGG GGT GAA ATT GTG CTG ACT CAG AGC CCA GGT ACC CTG TCT GTG TCT CCA
 25 GGC GAG AGA GCC ACC CTC TCC TGC CGG GCC AGT CAG AGC ATT GGT AGT AGC
 TTA CAC TGG TAC CAG CAG AAA CCA GGT CAG GCT CCA AGG CTT CTC ATC AAG
 30 TAT GCA TCC CAG TCC CTC TCA GGG ATC CCC GAT AGG TTC AGT GGC AGT GGA
 TCT GGG ACA GAT TTC ACC CTC ACC ATC AGT AGA CTG GAG CCT GAA GAT TTC
 GCA GTG TAT TAC TGT CAT CAG AGT AGT CGT TTA CCT CAC ACT TTC GGC CAA
 35 GGG ACC AAG GTG GAG ATC AAA CGT ACA

(SEQ ID NO: 78)

40 M S P S Q L I G F L L L W V P A S
 R G E I V L T Q S P G T L S V S P
 G E R A T L S C R A S Q S I G S S
 45 L H W Y Q Q K P G Q A P R L L I K
 Y A S Q S L S G I P D R F S G S G
 50 S G T D F T L T I S R L E P E D F
 A V Y Y C H Q S S R L P H T F G Q
 G T K V E I K R T

55 Modified 19D12/15H12 heavy chain-A (SEQ ID NO: 44)

39

5 ATG GAG TTT GGG CTG AGC TGG GTT TTC CTT GTT GCT ATA TTA AAA GGT GTC
 CAG TGT GAG GTT CAG CTG GTG CAG TCT GGG GGA GGC TTG GTA AAG CCT GGG
 GGG TCC CTG AGA CTC TCC TGT GCA GCC TCT GGA TTC ACC TTC AGT AGC TTT
 GCT ATG CAC TGG GTT CGC CAG GCT CCA GGA AAA GGT CTG GAG TGG ATA TCA
 10 GTT ATT GAT ACT CGT GGT GCC ACA TAC TAT GCA GAC TCC GTG AAG GGC CGA
 TTC ACC ATC TCC AGA GAC AAT GCC AAG AAC TCC TTG TAT CTT CAA ATG AAC
 AGC CTG AGA GCC GAG GAC ACT GCT GTG TAT TAC TGT GCA AGA CTG GGG AAC
 15 TTC TAC TAC GGT ATG GAC GTC TGG GGC CAA GGG ACC ACG GTC ACC GTC TCC
 TCA

(SEQ ID NO: 45)

20
 Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Lys Gly Val
 Gln Cys Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Lys Pro Gly
 25 Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Phe
 Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ser
 30 Val Ile Asp Thr Arg Gly Ala Thr Tyr Tyr Ala Asp Ser Val Lys Gly Arg
 Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu Gln Met Asn
 Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg Leu Gly Asn
 35 Phe Tyr Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser
 Ser

Modified 19D12/15H12 heavy chain-B (SEQ ID NO: 111)

40
 ATG GAG TTT GGG CTG AGC TGG GTT TTC CTT GTT GCT ATA TTA AAA GGT GTC
 CAG TGT GAG GTT CAG CTG GTG CAG TCT GGG GGA GGC TTG GTA CAG CCC GGG
 45 GGG TCC CTG AGA CTC TCC TGT GCA GCC TCT GGA TTC ACC TTC AGT AGC TTT
 GCT ATG CAC TGG GTT CGC CAG GCT CCA GGA AAA GGT CTG GAG TGG ATA TCA
 50 GTT ATT GAT ACT CGT GGT GCC ACA TAC TAT GCA GAC TCC GTG AAG GGC CGA
 TTC ACC ATC TCC AGA GAC AAT GCC AAG AAC TCC TTG TAT CTT CAA ATG AAC
 AGC CTG AGA GCC GAG GAC ACT GCT GTG TAT TAC TGT GCA AGA CTG GGG AAC
 55 TTC TAC TAC GGT ATG GAC GTC TGG GGC CAA GGG ACC ACG GTC ACC GTC TCC
 TCA

(SEQ ID NO: 112)

Met Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Lys Gly Val
5 Gln Cys Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Gln Pro Gly
Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Phe
Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ser
10 Val Ile Asp Thr Arg Gly Ala Thr Tyr Tyr Ala Asp Ser Val Lys Gly Arg
Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu Gln Met Asn
Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg Leu Gly Asn
15 Phe Tyr Tyr Gly Met Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser
Ser

20 **Gene Therapy**

The anti-IGFR1 antibodies of the invention may also be administered to a subject in a gene therapy approach. In a gene therapy approach, the cells of a subject are transformed with nucleic acids which encode the antibodies of the invention. Subjects comprising the nucleic acids will then produce the antibody molecules endogenously. Previously, Alvarez, *et al.*, (2000) (Clinical Cancer Research 6:3081-3087) introduced single-chain anti-ErbB2 antibodies to subjects using a gene therapy approach. The methods disclosed by Alvarez, *et al.*, may be easily adapted for the introduction of nucleic acids encoding an anti-IGFR1 antibody molecule of the invention to a subject.

30 Although nucleic acids encoding any polypeptide or antibody molecule of the invention may be introduced to a subject, in preferred embodiments, the antibody molecule is a fully human, single-chain antibody.

The nucleic acids may be introduced to the cells of a subject by any means known in the art. In preferred embodiments, the nucleic acids are introduced as part of a viral vector. Examples of preferred viruses from which the vectors may be derived include lentiviruses, herpes viruses, adenoviruses, adeno-associated viruses, vaccinia virus, baculovirus, alphavirus, influenza virus, and other recombinant viruses with desirable cellular tropism.

Various companies produce viral vectors commercially, including, but by no means limited to, Avigen, Inc. (Alameda, CA; AAV vectors), Cell Genesys (Foster City, CA; retroviral, adenoviral, AAV vectors, and lentiviral vectors), Clontech (retroviral and baculoviral vectors), Genovo, Inc. (Sharon Hill, PA; adenoviral and AAV vectors), Genvec (adenoviral vectors), IntroGene (Leiden, Netherlands;

adenoviral vectors), Molecular Medicine (retroviral, adenoviral, AAV, and herpes viral vectors), Norgen (adenoviral vectors), Oxford BioMedica (Oxford, United Kingdom; lentiviral vectors), and Transgene (Strasbourg, France; adenoviral, vaccinia, retroviral, and lentiviral vectors).

- 5 Methods for constructing and using viral vectors are known in the art (see, *e.g.*, Miller, *et al.*, (1992) *BioTechniques* 7:980-990). Preferably, the viral vectors are replication defective, that is, they are unable to replicate autonomously, and thus are not infectious, in the target cell. Preferably, the replication defective virus is a minimal virus, *i.e.*, it retains only the sequences of its genome which are necessary for
- 10 encapsidating the genome to produce viral particles. Defective viruses, which entirely or almost entirely lack viral genes, are preferred. Use of defective viral vectors allows for administration to cells in a specific, localized area, without concern that the vector can infect other cells. Thus, a specific tissue can be specifically targeted.

- Examples of vectors comprising attenuated or defective DNA virus sequences
- 15 include, but are not limited to, a defective herpes virus vector (Kanno, *et al.*, (1999) *Cancer Gen. Ther.* 6:147-154; Kaplitt, *et al.*, (1997) *J. Neurosci. Meth.* 71:125-132 and Kaplitt, *et al.*, (1994) *J. Neuro Onc.* 19:137-147).

- Adenoviruses are eukaryotic DNA viruses that can be modified to efficiently deliver a nucleic acid of the invention to a variety of cell types. Attenuated adenovirus
- 20 vectors, such as the vector described by Stratford-Perricaudet, *et al.*, (1992) (*J. Clin. Invest.* 90:626-630) are desirable in some instances. Various replication defective adenovirus and minimum adenovirus vectors have been described (PCT Publication Nos. WO94/26914, WO94/28938, WO94/28152, WO94/12649, WO95/02697 and WO96/22378). The replication defective recombinant adenoviruses according to the
- 25 invention can be prepared by any technique known to a person skilled in the art (Levrero, *et al.*, (1991) *Gene* 101:195; EP 185573; Graham, (1984) *EMBO J.* 3:2917; Graham, *et al.*, (1977) *J. Gen. Virol.* 36:59).

- The adeno-associated viruses (AAV) are DNA viruses of relatively small size which can integrate, in a stable and site-specific manner, into the genome of the cells
- 30 which they infect. They are able to infect a wide spectrum of cells without inducing any effects on cellular growth, morphology or differentiation, and they do not appear to be involved in human pathologies. The use of vectors derived from the AAVs for transferring genes *in vitro* and *in vivo* has been described (see Daly, *et al.*, (2001) *Gene Ther.* 8:1343-1346, 1245-1315; Larson, *et al.*, (2001) *Adv. Exp. Med. Bio.*

489:45-57; PCT Publication Nos. WO91/18088 and WO93/09239; U.S. Patent Nos. 4,797,368 and 5,139,941 and EP 488528B1).

In another embodiment, the gene can be introduced in a retroviral vector, *e.g.*, as described in U.S. Patent Nos. 5,399,346, 4,650,764, 4,980,289, and 5,124,263; 5 Mann, *et al.*, (1983) Cell 33:153; Markowitz, *et al.*, (1988) J. Virol., 62:1120; EP 453242 and EP178220. The retroviruses are integrating viruses which infect dividing cells.

Lentiviral vectors can be used as agents for the direct delivery and sustained expression of nucleic acids encoding an antibody molecule of the invention in several 10 tissue types, including brain, retina, muscle, liver and blood. The vectors can efficiently transduce dividing and nondividing cells in these tissues, and maintain long-term expression of the antibody molecule. For a review, see Zufferey, *et al.*, (1998) J. Virol. 72:9873-80 and Kafri, *et al.*, (2001) Curr. Opin. Mol. Ther. 3:316-326. Lentiviral packaging cell lines are available and known generally in the art. They 15 facilitate the production of high-titer lentivirus vectors for gene therapy. An example is a tetracycline-inducible VSV-G pseudotyped lentivirus packaging cell line which can generate virus particles at titers greater than 10⁶ IU/ml for at least 3 to 4 days; see Kafri, *et al.*, (1999) (J. Virol. 73: 576-584). The vector produced by the inducible cell line can be concentrated as needed for efficiently transducing nondividing cells *in* 20 *vitro* and *in vivo*.

Sindbis virus is a member of the alphavirus genus and has been studied extensively since its discovery in various parts of the world beginning in 1953. Gene transduction based on alphavirus, particularly Sindbis virus, has been well-studied *in vitro* (see Straus, *et al.*, (1994) Microbiol. Rev., 58:491-562; Bredenbeek, *et al.*, 25 (1993) J. Virol., 67: 6439-6446 Iijima, *et al.*, (1999) Int. J. Cancer 80:110-118 and Sawai, *et al.*, (1998) Biochim. Biophys. Res. Comm. 248:315-323). Many properties of alphavirus vectors make them a desirable alternative to other virus-derived vector systems being developed, including rapid engineering of expression constructs, production of high-titered stocks of infectious particles, infection of nondividing cells, 30 and high levels of expression (Strauss, *et al.*, (1994) Microbiol. Rev. 58:491-562). Use of Sindbis virus for gene therapy has been described. (Wahlfors, *et al.*, (2000) Gene Ther. 7:472-480 and Lundstrom (1999) J. Recep. Sig. Transduct. Res. 19(1-4):673-686).

In another embodiment, a vector can be introduced to cells by lipofection or with other transfection facilitating agents (peptides, polymers, *etc.*). Synthetic cationic lipids can be used to prepare liposomes for *in vivo* and *in vitro* transfection of a gene encoding a marker (Felgner, *et al.*, (1987) Proc. Natl. Acad. Sci. USA 84:7413-7417 and Wang, *et al.*, (1987) Proc. Natl. Acad. Sci. USA 84:7851-7855). Useful lipid compounds and compositions for transfer of nucleic acids are described in PCT Publication Nos. WO 95/18863 and WO96/17823, and in U.S. Patent No. 5,459,127.

It is also possible to introduce the vector *in vivo* as a naked DNA plasmid. Naked DNA vectors for gene therapy can be introduced into the desired host cells by methods known in the art, *e.g.*, electroporation, microinjection, cell fusion, DEAE dextran, calcium phosphate precipitation, use of a gene gun, or use of a DNA vector transporter (see, *e.g.*, Wilson, *et al.*, (1992) J. Biol. Chem. 267:963-967; Williams, *et al.*, (1991) Proc. Natl. Acad. Sci. USA 88:2726-2730). Receptor-mediated DNA delivery approaches can also be used (Wu, *et al.*, (1988) J. Biol. Chem. 263:14621-14624). U.S. Patent Nos. 5,580,859 and 5,589,466 disclose delivery of exogenous DNA sequences, free of transfection facilitating agents, in a mammal. Recently, a relatively low voltage, high efficiency *in vivo* DNA transfer technique, termed electrotransfer, has been described (Vilquin, *et al.*, (2001) Gene Ther. 8:1097; Payen, *et al.*, (2001) Exp. Hematol. 29:295-300; Mir (2001) Bioelectrochemistry 53:1-10; PCT Publication Nos. WO99/01157, WO99/01158 and WO99/01175).

Pharmaceutical Compositions

An antibody or antigen-binding fragment of the invention can be incorporated into a pharmaceutical composition, along with a pharmaceutically acceptable carrier, suitable for administration to a subject *in vivo*. Although the scope of the present invention includes pharmaceutical compositions which may be administered to a subject by any route (*e.g.*, oral, ocular, topical or pulmonary (inhalation)), administration by a parenteral route such as intratumoral injection, intravenous injection, subcutaneous injection or intramuscular injection is preferred. In a preferred embodiment, the pharmaceutical compositions of the invention comprise 1H3, 15H12, 19D12, 15H12/19D12 LCA, 15H12/19D12 LCB, 15H12/19D12 LCC, 15H12/19D12 LCD, 15H12/19D12 LCE, 15H12/19D12 LCF, 15H12/19D12 HCA or 15H12/19D12 HCB and a pharmaceutically acceptable carrier.

- For general information concerning formulations, see, *e.g.*, Gilman, *et al.*, (eds.) (1990), The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; A. Gennaro (ed.), Remington's Pharmaceutical Sciences, 18th Edition, (1990), Mack Publishing Co., Easton, Pennsylvania.; Avis, *et al.*, (eds.) (1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, New York; Lieberman, *et al.*, (eds.) (1990) Pharmaceutical Dosage Forms: Tablets Dekker, New York; and Lieberman, *et al.*, (eds.) (1990), Pharmaceutical Dosage Forms: Disperse Systems Dekker, New York, Kenneth A. Walters (ed.) (2002) Dermatological and Transdermal Formulations (Drugs and the Pharmaceutical Sciences), Vol 119, Marcel Dekker.
- 10 Pharmaceutically acceptable carriers are conventional and very well known in the art. Examples include aqueous and nonaqueous carriers, stabilizers, antioxidants, solvents, dispersion media, coatings, antimicrobial agents, buffers, serum proteins, isotonic and absorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for injection into a
- 15 subject's body.
- Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic
- 20 esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.
- Stabilizers, such as α , α -trehalose dihydrate may be included for stabilizing the antibody molecules of the invention from degrading effects of dessication or freeze-
- 25 drying.
- Examples of pharmaceutically-acceptable antioxidants include: water soluble antioxidants such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; and oil-soluble antioxidants such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT),
- 30 lecithin, propyl gallate, alpha-tocopherol, and the like; and metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

Prevention of the presence of microorganisms may be ensured both by sterilization procedures, and by the inclusion of various antimicrobial agents such as EDTA, EGTA, paraben, chlorobutanol, phenol sorbic acid, and the like.

Suitable buffers which may be included in the pharmaceutical compositions of the invention include L-histidine based buffers, phosphate based buffers (*e.g.*,
5 phosphate buffered saline, pH \approx 7), sorbate based buffers or glycine-based buffers.

Serum proteins which may be included in the pharmaceutical compositions of the invention may include human serum albumin.

Isotonic agents, such as sugars, ethanol, polyalcohols (*e.g.*, glycerol,
10 propylene glycol, liquid polyethylene glycol, mannitol or sorbitol), sodium citrate or sodium chloride (*e.g.*, buffered saline) may also be included in the pharmaceutical compositions of the invention.

Prolonged absorption of an injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum
15 monostearate and/or gelatin.

Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils.

Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile
20 injectable solutions or dispersions. The use of such media and agents for pharmaceutically active substances is well known in the art.

Sterile injectable solutions can be prepared by incorporating the antibody or antigen-binding fragment of the invention in the required amount in an appropriate solvent, optionally with one or a combination of ingredients enumerated above, as
25 required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the antibody molecule into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying (lyophilization)
30 that yield a powder of the active ingredient plus any additional, desired ingredient from a previously sterile-filtered solution thereof.

The antibody or antigen-binding fragment of the invention may also be orally administered. Pharmaceutical compositions for oral administration may contain, in addition to the binding composition, additives such as starch (*e.g.*, potato, maize or

wheat starch or cellulose), starch derivatives (e.g., microcrystalline cellulose or silica), sugars (e.g., lactose), talc, stearate, magnesium carbonate or calcium phosphate. In order to ensure that oral compositions comprising an antibody or antigen-binding fragment of the invention are well tolerated by the patient's digestive system, mucus
5 formers or resins may be included. It may also be desirable to improve tolerance by formulating the antibody or antigen-binding fragment in a capsule which is insoluble in the gastric juices. An exemplary pharmaceutical composition of this invention in the form of a capsule is prepared by filling a standard two-piece hard gelatin capsule with the antibody or antigen-binding fragment of the invention in powdered form, lactose,
10 talc and magnesium stearate. Oral administration of immunoglobulins has been described (Foster, *et al.*, (2001) Cochrane Database System rev. 3:CD001816)

An antibody or antigen-binding fragment of the invention may also be included in a pharmaceutical composition for topical administration. Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration
15 through the skin to the site where treatment is required, such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose.

Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the antibody or antigen-
20 binding fragment in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile, aqueous solution
25 optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid
30 formulations of the active ingredient for external application. They may be made by mixing the antibody or antigen-binding fragment of the invention in finely-divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy basis. The basis may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a

metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol or macrogels. The formulation may incorporate any suitable surface active agent such as an anionic, cationic or non-
5 ionic surface active such as sorbitan esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

The antibodies and antigen-binding fragments of the invention may also be
10 administered by inhalation. A suitable pharmaceutical composition for inhalation may be an aerosol. An exemplary pharmaceutical composition for inhalation of an antibody or antigen-binding fragment of the invention may include: an aerosol container with a capacity of 15-20 ml comprising the antibody or antigen-binding fragment of the invention, a lubricating agent, such as polysorbate 85 or oleic acid,
15 dispersed in a propellant, such as freon, preferably in a combination of 1,2-dichlorotetrafluoroethane and difluorochloromethane. Preferably, the composition is in an appropriate aerosol container adapted for either intranasal or oral inhalation administration.

In yet another embodiment of the present invention, the pharmaceutical
20 composition can be administered by combination therapy. For example, the combination therapy can include a pharmaceutical composition of the present invention in association with one or more anti-cancer therapeutic agents (*e.g.*, alkylating agents, antimetabolites, anti-tumor antibiotics, mitotic inhibitors, chromatin function inhibitors, anti-angiogenesis agents, anti-estrogens, anti-androgens, antibody
25 therapies or immunomodulators). An "anti-cancer therapeutic agent" is a substance which, when administered to a subject, treats or prevents the development of cancer in the subject's body. The compositions of the invention may be administered in association with one or more anti-cancer therapeutic procedures (*e.g.*, radiation therapy or surgical tumorectomy). An "anti-cancer therapeutic procedure" is a
30 process which is performed on a subject which treats or reduces the incidence of cancer in the subject. When a combination therapy is used, the antibodies or antigen-binding fragments of the invention, or pharmaceutical compositions thereof, may be formulated into a single composition for simultaneous delivery or formulated separately into two or more compositions (*e.g.*, a kit). Furthermore, the antibody or

antigen-binding fragment may be administered to a subject at a different time than when the other therapeutic agent or therapeutic procedure is administered; for example, each administration may be given non-simultaneously at several intervals over a given period of time.

5 "Alkylating agent" refers to any substance which can cross-link or alkylate any molecule, preferably nucleic acid (e.g., DNA), within a cell. Examples of alkylating agents include mechlorethamine, cyclophosphamide, ifosfamide, phenylalanine mustard, melphalen, chlorambucol, uracil mustard, estramustine, thiotepa, busulfan, lomustine, carmustine, streptozocin, dacarbazine, cis-platinum, carboplatin and
10 altretamine.

"Antimetabolites" refer to substances that block cell growth and/or metabolism by interfering with certain activities, usually DNA synthesis. Examples of antimetabolites include methotrexate, 5-fluorouracil, floxuridine, 5-fluorodeoxyuridine, capecitabine, fludarabine, cytosine arabinoside, 6-mercaptopurine, 6-thioguanine,
15 gemcitabine, cladribine, deoxycoformycin and pentostatin.

"Anti-tumor antibiotics" refer to compounds which may prevent or inhibit DNA, RNA and/or protein synthesis. Examples of anti-tumor antibiotics include doxorubicin, daunorubicin, idarubicin, valrubicin, mitoxantrone, dactinomycin, mithramycin, plicamycin, mitomycin C, bleomycin, and procarbazine.

20 "Mitotic inhibitors" prevent normal progression of the cell cycle and mitosis. In general, microtubule inhibitors such as paclitaxel and docetaxel are capable of inhibiting mitosis. Vinca alkaloids such as vinblastine, vincristine and vinorelbine are also capable of inhibiting mitosis.

"Chromatin function inhibitors" refer to substances which inhibit the normal
25 function of chromatin modeling proteins such as topoisomerase I or topoisomerase II. Examples of chromatin function inhibitors include topotecan, irinotecan, etoposide and teniposide.

"Anti-angiogenesis agent" refers to any drug, compound, substance or agent which inhibits growth of blood vessels. Exemplary anti-angiogenesis agents include,
30 but are by no means limited to, razoxin, marimastat, COL-3, neovastat, BMS-275291, thalidomide, squalamine, endostatin, SU5416, SU6668, interferon-alpha, EMD121974, interleukin-12, IM862, angiostatin and vitaxin.

"Anti-estrogen" or "anti-estrogenic agent" refer to any substance which reduces, antagonizes or inhibits the action of estrogen. Examples of anti-estrogen

agents are tamoxifen, toremifene, raloxifene, droloxifene, idoxifene, anastrozole, letrozole, and exemestane.

"Anti-androgens" or "anti-androgen agents" refer to any substance which reduces, antagonizes or inhibits the action of an androgen. Examples of anti-androgens are flutamide, nilutamide, bicalutamide, spironolactone, cyproterone acetate, finasteride and cimitidine.

Antibody therapies which may be administered in conjunction with the antibodies or antigen-binding fragments of the invention include trastuzumab (*e.g.*, herceptin) (see, for example, Sliwkowski, *et al.*, (1999) *Semin. Oncol.* 26(4 Suppl 12):60-70), vitaxin and rituximab.

"Immunomodulators" are substances which stimulate the immune system. Examples of immunomodulators include denileukin diftitox, levamisole in conjunction with 5-fluorouracil, interferon and interleukin-2.

"Radiotherapy" or "radiation therapy" refers to treating a disease, such as cancer, by administration of ionizing radiation (preferably to a tumor site). Examples of ionizing radiation which may be administered include X-rays, gamma rays (*e.g.*, emitted by radium, uranium or cobalt 60), and particle beam radiation (*e.g.*, protons, neutrons, pions or heavy ions).

20

Dosage

Preferably, an antibody or antigen-binding fragment of the invention is administered to a subject at a "therapeutically effective dosage" which preferably inhibits a disease or condition which is mediated by IGFR1 (*e.g.*, tumor growth) to any extent-preferably by at least about 20%, more preferably by at least about 40%, even more preferably by at least about 60%, and still more preferably by at least about 80%-100% relative to untreated subjects. The ability of an antibody or antigen-binding fragment of the invention to inhibit cancer can be evaluated in an animal model system predictive of efficacy in human tumors. Alternatively, this property of a composition can be evaluated by examining the ability of an antibody or antigen-binding fragment of the invention to inhibit tumor cell growth *in vitro* by assays (see below) well-known to the skilled practitioner. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

Dosage regimens may be adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage.

A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the antibody or antigen-binding fragment of the invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, a suitable daily dose of a composition of the invention may be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. It is preferred that administration be by injection, preferably proximal to the site of the target (e.g., tumor). If desired, the effective daily dose of a pharmaceutical composition may be administered as two, three, four, five, six or more subdoses administered separately at appropriate intervals throughout the day.

Therapeutic Methods and Administration

The antibodies or antigen-binding fragments of the invention and pharmaceutical compositions comprising the antibodies or antigen-binding fragments of the invention may be used for treating or preventing any disease or condition in a subject which is mediated by elevated expression or activity of IGFR1 or by elevated expression of its ligand (e.g., IGF-I or IGF-II) and which may be treated or prevented by modulation of IGFR1 ligand binding, activity or expression. Preferably, the disease or condition is mediated by an increased level of IGFR1, IGF-I or IGF-II and is treated or prevented by decreasing IGFR1 ligand binding, activity (e.g., autophosphorylation activity) or expression. Preferably, the disease or condition is malignancy, more preferably a malignancy characterized by a tumor which expresses IGFR1, such as, but not limited to, bladder cancer, Wilm's cancer, bone cancer, prostate cancer, lung cancer, colorectal cancer, breast cancer, cervical cancer, synovial sarcoma, ovarian cancer, pancreatic cancer, benign prostatic hyperplasia (BPH), diarrhea associated

with metastatic carcinoid and vasoactive intestinal peptide secreting tumors (*e.g.*, VIPoma or Werner-Morrison syndrome). Acromegaly may also be treated with the antibody molecules of the invention. Antagonism of IGF-I has been reported for treatment of acromegaly (Drake, *et al.*, (2001) Trends Endocrin. Metab. 12: 408-413).

- 5 Other non-malignant medical conditions which may also be treated, in a subject, by administering an anti-IGFR1 antibody of the invention include gigantism, psoriasis, atherosclerosis, smooth muscle restenosis of blood vessels or inappropriate microvascular proliferation, such as that found as a complication of diabetes, especially of the eye.

- 10 The term "subject" may refer to any organism, preferably an animal, more preferably a mammal (*e.g.*, rat, mouse, dog, cat, rabbit) and most preferably a human.

- In preferred embodiments, the antibodies and antigen-binding fragments of the invention and pharmaceutical compositions thereof are administered by an invasive route such as by injection (see above). Administration by a non-invasive route (see above) is also within the scope of the present invention.

Compositions can be administered with medical devices known in the art. For example, in a preferred embodiment, a pharmaceutical composition of the invention can be administered by injection with a hypodermic needle.

- 20 The pharmaceutical compositions of the invention may also be administered with a needleless hypodermic injection device; such as the devices disclosed in U.S. Patent Nos. 5,399,163; 5,383,851; 5,312,335; 5,064,413; 4,941,880; 4,790,824 or 4,596,556.

- Examples of well-known implants and modules useful in the present invention include: U.S. Patent No. 4,487,603, which discloses an implantable micro-infusion pump for dispensing medication at a controlled rate; U.S. Patent No. 4,447,233, which discloses a medication infusion pump for delivering medication at a precise infusion rate; U.S. Patent No. 4,447,224, which discloses a variable flow implantable infusion apparatus for continuous drug delivery; U.S. Patent. No. 4,439,196, which discloses an osmotic drug delivery system having multi-chamber compartments.
- 30 Many other such implants, delivery systems, and modules are well known to those skilled in the art.

Assays

The anti-IGFR1 antibodies may be used to detect IGFR1 in a biological sample *in vitro* or *in vivo* (see, for example, Zola, Monoclonal Antibodies: A Manual of Techniques, pp. 147-158 (CRC Press, Inc., 1987)). The anti-IGFR1 antibodies may
5 be used in a conventional immunoassay, including, without limitation, an ELISA, an RIA, FACS, tissue immunohistochemistry, Western blot or immunoprecipitation. The anti-IGFR1 antibodies of the invention may be used to detect IGFR1 from humans. The invention provides a method for detecting IGFR1 in the biological sample comprising contacting the biological sample with an anti-IGFR1 antibody of the
10 invention and detecting the anti-IGFR1 antibody bound to IGFR1, thereby indicating the presence of the IGFR1 in the biological sample. In one embodiment, the anti-IGFR1 antibody is directly labeled with a detectable label and may be detected directly. In another embodiment, the anti-IGFR1 antibody (the first antibody) is unlabeled and a secondary antibody or other molecule that can bind the anti-IGFR1
15 antibody is labeled. As is well known to one of skill in the art, a secondary antibody is chosen that is able to specifically bind the specific species and class of the first antibody. For example, if the anti-IGFR1 antibody is a human IgG, then the secondary antibody may be an anti-human-IgG. The presence of an anti-IGFR1/IGFR1 complex in the biological sample can be detected by detecting the
20 presence of the labeled secondary antibody. Other molecules that can bind to antibodies (*e.g.*, anti-IGFR1 antibodies) include, without limitation, Protein A and Protein G, both of which are available commercially, *e.g.*, from Pierce Chemical Co. (Rockford, IL)

Suitable labels for the anti-IGFR1 antibody or secondary antibody have been
25 disclosed *supra*, and include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, magnetic agents and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable
30 fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; an example of a magnetic agent includes gadolinium; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

In an alternative embodiment, IGFR1 can be assayed in a biological sample by a competition immunoassay utilizing IGFR1 standards labeled with a detectable substance and an unlabeled anti-IGFR1 antibody. In this assay, the biological sample, the labeled IGFR1 standards and the anti-IGFR1 antibody are combined and the amount of labeled IGFR1 standard bound to the unlabeled antibody is
5 determined. The amount of IGFR1 in the biological sample is inversely proportional to the amount of labeled IGFR1 standard bound to the anti-IGFR1 antibody. One may use the immunoassays disclosed above for a number of purposes. In one embodiment, the anti-IGFR1 antibodies may be used to detect IGFR1 in cells in cell
10 culture. In a preferred embodiment, the anti-IGFR1 antibodies may be used to determine the level of tyrosine phosphorylation, tyrosine autophosphorylation of IGFR1, and/or the amount of IGFR1 on the cell surface after treatment of the cells with various compounds. This method can be used to test compounds that may be used to activate or inhibit IGFR1. In this method, one sample of cells is treated with a
15 test compound for a period of time while another sample is left untreated. If tyrosine autophosphorylation is to be measured, the cells are lysed and tyrosine phosphorylation of the IGFR1 is measured using an immunoassay, for example, as described above. If the total level of IGFR1 is to be measured, the cells are lysed and the total IGFR1 level is measured using one of the immunoassays described
20 above.

A preferred immunoassay for determining IGFR1 tyrosine phosphorylation or for measuring total IGFR1 levels is an ELISA or Western blot. If only the cell surface level of IGFR1 is to be measured, the cells are not lysed, and the cell surface levels of IGFR1 are measured using one of the immunoassays described above. A
25 preferred immunoassay for determining cell surface levels of IGFR1 includes the steps of labeling the cell surface proteins with a detectable label, such as biotin or ¹²⁵I, immunoprecipitating the IGFR1 with an anti-IGFR1 antibody and then detecting the labeled IGFR1. Another preferred immunoassay for determining the localization of IGFR1, *e.g.*, cell surface levels, is by using immunohistochemistry. Methods such as
30 ELISA, RIA, Western blot, immunohistochemistry, cell surface labeling of integral membrane proteins and immunoprecipitation are well known in the art. In addition, the immunoassays may be scaled up for high throughput screening in order to test a large number of compounds for either activation or inhibition of IGFR1.

The anti-IGFR1 antibodies of the invention may also be used to determine the levels of IGFR1 in a tissue or in cells derived from the tissue. In a preferred embodiment, the tissue is a diseased tissue. In a more preferred embodiment, the tissue is a tumor or a biopsy thereof. In a preferred embodiment of the method, a tissue or a biopsy thereof is excised from a patient. The tissue or biopsy is then used in an immunoassay to determine, *e.g.*, IGFR1 levels, cell surface levels of IGFR1, levels of tyrosine phosphorylation of IGFR1, or localization of IGFR1 by the methods discussed above. The method can be used to determine if a tumor expresses IGFR1 at a high level.

The above-described diagnostic method can be used to determine whether a tumor expresses high levels of IGFR1, which may be indicative that the tumor will respond well to treatment with anti-IGFR1 antibody. The diagnostic method may also be used to determine whether a tumor is potentially cancerous, if it expresses high levels of IGFR1, or benign, if it expresses low levels of IGFR1. Further, the diagnostic method may also be used to determine whether treatment with anti-IGFR1 antibody is causing a tumor to express lower levels of IGFR1 and/or to exhibit lower levels of tyrosine autophosphorylation, and thus can be used to determine whether the treatment is successful. In general, a method to determine whether an anti-IGFR1 antibody decreases tyrosine phosphorylation comprises the steps of measuring the level of tyrosine phosphorylation in a cell or tissue of interest, incubating the cell or tissue with an anti-IGFR1 antibody or antigen-binding portion thereof, then re-measuring the level of tyrosine phosphorylation in the cell or tissue. The tyrosine phosphorylation of IGFR1 or of another protein(s) may be measured. The diagnostic method may also be used to determine whether a tissue or cell is not expressing high enough levels of IGFR1 or high enough levels of activated IGFR1, which may be the case for individuals with dwarfism, osteoporosis or diabetes. A diagnosis that levels of IGFR1 or active IGFR1 are too low could be used for treatment with activating anti-IGFR1 antibodies, IGF-1, IGF-2 or other therapeutic agents for increasing IGFR1 levels or activity.

The antibodies of the present invention may also be used *in vivo* to localize tissues and organs that express IGFR1. In a preferred embodiment, the anti-IGFR1 antibodies can be used to localize IGFR1-expressing tumors. The advantage of the anti-IGFR1 antibodies of the present invention is that they will not generate an immune response upon administration. The method comprises the steps of

administering an anti-IGFR1 antibody or a pharmaceutical composition thereof to a patient in need of such a diagnostic test and subjecting the patient to imaging analysis to determine the location of the IGFR1-expressing tissues. Imaging analysis is well known in the medical art, and includes, without limitation, x-ray analysis, magnetic resonance imaging (MRI) or computed tomography (CT). In another embodiment of the method, a biopsy is obtained from the patient to determine whether the tissue of interest expresses IGFR1 rather than subjecting the patient to imaging analysis. In a preferred embodiment, the anti-IGFR1 antibodies may be labeled with a detectable agent that can be imaged in a patient. For example, the antibody may be labeled with a contrast agent, such as barium, which can be used for x-ray analysis, or a magnetic contrast agent, such as a gadolinium chelate, which can be used for MRI or CE. Other labeling agents include, without limitation, radioisotopes, such as ⁹⁹Tc. In another embodiment, the anti-IGFR1 antibody will be unlabeled and will be imaged by administering a secondary antibody or other molecule that is detectable and that can bind the anti-IGFR1 antibody.

EXAMPLES

The following examples are provided to further describe the present invention and should not be construed to limit the scope of the invention in any way.

EXAMPLE 1: Construction of Fully Human Anti-IGFR1 Antibodies.

1.0. Introduction.

Fully human monoclonal antibodies specific for human insulin-like growth factor receptor 1 (IGFR1) were generated from HuMab mice of the Hco7 genotype (see below), immunized with recombinant sIGFR1 and IGFR1 transfected HEK293 cells. A detailed description of Hco7 mice is provided in U.S. Patent Nos. 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016; 5,770,429; 5,789,650; 5,814,318; 5,874,299 and 5,877,397 and in Harding, *et al.*, (1995) Ann. NY Acad. Sci. 764:536-546. Antibodies 1H3, 15H12 and 19D12 were isolated from a HuMab mouse (referred to herein as #23716) which was selected for fusion based on the presence of antigen specific serum IgG titers of 25,600 to the immunizing antigen. The 1H3, 15H12 and 19D12 antibodies were found to bind IGFR1.

2.0 Materials and Methods and Results.

2.1. Antigen.

5

2.1.1. Mice were immunized with two forms of antigen: (1) live cells (*IGFR1* transfected HEK293 cells) and (2) purified protein (sIGFR1; an NSO-expressed recombinant protein encompassing the α -subunit and the extracellular domain of the β -subunit of IGFR1). The biologically active version of this protein is in glycosylated form.

10

2.1.2. Three immunizations with soluble IGFR1 antigen and final tail vein boosts were performed with a purified IGFR1 preparation at a concentration of 2.67 mg/ml. Soluble IGFR1 was mixed with either complete or incomplete Freund's adjuvant (CFA and IFA) and mice were injected with 0.2 cc (cubic centimeters) prepared antigen into the intraperitoneal cavity. Final tail vein immunizations were performed with soluble IGFR1 in sterile PBS (phosphate buffer saline).

15

2.1.3. Immunizations were also performed with HEK293 cells transfected with *IGFR1* DNA. Specifically, each mouse was immunized, by injection into the intraperitoneal cavity, with 0.2 cc of sterile saline containing $1.0\text{-}2.0 \times 10^7$ HEK293 cells expressing IGFR1.

20

2.2. Transgenic Mice.

25

2.2.1. Mice were housed in filter cages and were evaluated to be in good physical condition at the time of immunization, at the time of the bleeds and on the day fusions were produced.

30

2.2.2. The mouse that produced the selected hybridomas was a male (mouse ID #23716) of the (CMD)++; (Hco7) 11952+; (JKD) ++; (KCo5) 9272+ genotype. Individual transgene designations are in parentheses, followed by line numbers for randomly integrated transgenes. The symbols ++ and + indicate homozygous or hemizygous; however, because the mice are routinely screened using a PCR-based assay that does not allow us to distinguish between heterozygosity and homozygosity for the randomly integrated human Ig transgenes, a +

35

designation may be given to mice that are actually homozygous for these elements.

2.3. Immunization Procedure and Schedule.

5

2.3.1. The immunization schedule is shown in the following table.

Table 2. Mouse immunization schedule.

Day	Immunization: adjuvant, antigen	Bleed and Titer ¹
Day 1	1.0 X 10 ⁷ live IGFR1 transfected HEK293 cells in saline	
Day 15	CFA adjuvant, sIGFR1 (20µg)	
Day 29	1.0 X 10 ⁷ live IGFR1 transfected HEK293 cells in saline	
Day 37		antibody titer measured
Day 43	IFA adjuvant, sIGFR1 (~ 40 µg)	
Day 54		antibody titer measured
Day 57	1.0 X 10 ⁷ live IGFR1 transfected HEK293 cells in saline	
Day 96	1.0 X 10 ⁷ live IGFR1 transfected HEK293 cells in saline	
Day 103		antibody titer measured
Day 112	CFA adjuvant, sIGFR1 (25 µg)	
Day 126		antibody titer measured
Days 128 and 129	Final tail vein intravenous boosts with sIGFR1 ²	

10

¹ Titer information is shown below.

² Fusions were performed on day 131.

15 **Table 3. Titers of IGFR1 specific antibody during the immunization period of mouse 23716 described in Table 2 (see above).**

Day	Titer
37	100

54	800
103	6400
126	25600

2.4. Hybridoma Preparation and Testing.

5 2.4.1. The SP2/0-AG14 myeloma cell line (ATCC CRL 1581) was used for the fusions. The original ATCC vial was thawed and expanded in culture. A seed stock of frozen vials was prepared from this expansion. Cells were maintained in culture for 6-8 weeks and passed twice a week.

10 2.4.2. High Glucose DMEM containing 10% FBS, Antibiotic-antimycotic (100X), and 0.1% L-glutamine was used to culture myeloma cells. Additional media supplements were added to the hybridoma growth media which included: 5% Origen - Hybridoma Cloning Factor (Fischer Scientific; Suwanee, GA), 4.5×10^{-4} M sodium Pyruvate, HAT 1.0×10^{-4} M Hypoxanthine, 4.0×10^{-7} M Aminopterin, 1.6×10^{-5} M Thymidine, or
15 HT 1.0×10^{-4} M Hypoxanthine, 1.6×10^{-5} M Thymidine; and characterized fetal bovine serum.

2.4.3. The spleen from mouse number #23716 was normal in size and yielded 5.73×10^8 viable cells.

20 2.4.4. The splenocytes were fused according to the following procedure:

1. Place approximately 10 ml of DMEM + 10%FBS into a 50 mL tube.
2. Sacrifice the intravenously boosted mouse.
3. Transfer the mouse into a hood onto a paper towel.
- 25 4. Soak the mouse with alcohol, and place onto its right side - left side up.
5. Make a small cut into the skin above the spleen area.
6. Pull skin away from the mouse using both hands.
7. Soak with alcohol again.
- 30 8. Use sterile instruments to open the peritoneum.

9. Insert the scissor points under the spleen and open the scissors so as to allow room to grasp the spleen with the forceps.
10. Remove the spleen and place into the tube containing DMEM + 10%FBS. Transfer to a sterile tissue culture room.
- 5 11. Inside a sterile hood, add approximately 7 mL of DMEM without serum to each of 2 sterile 60 mm culture dishes.
12. Transfer the spleen to the first dish.
13. Remove any adhesions from the spleen using sterile instruments.
- 10 14. Place a sterile homogenizer base into a test tube rack (for support).
15. Add the cleaned spleen into the homogenizer.
16. Add approximately 5 mL of DMEM and homogenize 4 passes. Pour off into a sterile 50 mL centrifuge tube.
- 15 17. Add another 5-6 mL of DMEM into the homogenizer and make another 3-4 additional passes.
18. Pour off into the same tube as described above.
19. Spin the cells at 1000 rpm for 10 minutes in a centrifuge.
20. Remove the supernatant. Pour off and resuspend pellets in
20 DMEM.
21. Count the spleen cells.
22. Transfer appropriate volume of SP2/0 cells (6 spleen cells per 1 cell of SP2/0) to a 50 mL centrifuge tube. Record volume.
23. Adjust volume of spleen cells with DMEM for more
25 convenient balancing for centrifugation.
24. Spin cells for 10 minutes at 1000 rpm in a centrifuge.
25. Remove supernatants - pour off and resuspend pellets in 30-40 mL of DMEM wash medium (serum free). Combine all cells in one tube.
- 30 26. Spin again as above.
27. Pour off supernatant and resuspend pellet.
28. Add approximately 1.2 mL of PEG (polyethylene glycol) at about 1 minute while gently swirling the tube in a beaker containing 37°C water.

29. Let the tube sit for 90 seconds, then add 15 mL of DMEM wash medium at 3 minutes.

30. Spin the tube as described above.

31. Remove the supernatant and resuspend the pellet gently.

5 32. Add approximately 10 mL of Hat medium to the tube.

33. Pipette cells into the full volume of HAT medium. Allow the cells to sit for 30-60 minutes in an incubator before plating.

34. Plate cells into 96 well culture plates, 200 μ L/well (about 1×10^7 cells per 96-well plate).

10 35. Feed cells on day 7 with HT media, 250 μ L/well. (HT media, same as HAT media, with Aminopterin removed)

2.4.5. An initial ELISA screen for human IgG κ antibodies was performed 7-10 days post fusion according to the following procedure:

15 1. Coat plate overnight with anti-hu- κ , 1 μ g/mL or anti-hu- γ , 1 μ g/mL in 1X PBS, 50 μ L/well. Store in refrigerator.

2. Empty plate and block plate in 1X PBST (PBS with Tween) + 5% chicken serum for 1 hour at room temperature (100 μ L/well).

20 3. Empty plate and wash manually with wash bottle (3X) or plate washer (3X) using 1X PBST. If wash bottle used, drain plates on paper towels.

4. Standards are used for testing production level of the clones. Make dilutions with unknowns (1:10 in first well and dilute 2 fold across plate). Hu-IgG standards start at 1000 ng/mL and dilute 2 fold across plate. Leave a few wells for blanks: 1X PBST + 5% chicken serum which is used for dilutions, 100 μ L/well. Incubate at room temperature for 1 hour. Fusion screens and subclones are generally tested diluted 1:2 in blocking buffer. A positive control may also be used when screening fusions and subclones.

25 5. Repeat wash step #3.

30 6. Dilute secondary antibody HRP(horse radish peroxidase)-anti-hu IgG-Fc reagent 1:5000 or HRP-anti-hu- κ in 1XPBST +5% chicken serum, add 100 μ L/well. Incubate 1 hour at room temperature.

7. Repeat wash step#3. (2X)

8. Develop plate using 10 ml citrate phosphate buffer pH4.0, 80 μ L ABTS, 8 μ L H₂O₂ per plate.

9. Incubate 30 minutes to 1 hour at room temperature. Read plate at OD_{415 nm-490 nm}.

5

Solutions:

1X PBST= 1xPBS + 0.05% tween-20

Citrate phosphate buffer = 21 gm/L citric acid, 14.2 gm/L disodium hydrogen phosphate (anhydrous); pH4.0

10

ABTS= 27.8 mg/mL 2,2'-azino-bis (3-ethylbenz-thiazoline-6-sulfonic acid) diammonium salt in citrate buffer, freeze 1 mL aliquots.

Plate = 96 well assay plate.

A positive ELISA signal was detected in the wells corresponding to hybridomas 1H3, 15H12 and 19D12, demonstrating that these hybridomas produced human IgG antibodies.

15

2.4.6. Hybridoma supernatants corresponding to human IgG κ positive wells were then screened on soluble IGFR1 coated ELISA plates according to the following procedure:

20

1. Coat plate overnight with IGFR1 (1.0 μ g/mL) in 1X PBS, 50 μ L/well. Store in refrigerator. Five milliliters needed for coating plate.

2. Empty plate and block plate in 1X PBST + 5% chicken serum for 1 hour at room temperature (100 μ L/well).

25

3. Empty plate and wash manually with wash bottle (3X) or plate washer (3X) using 1X PBST. If wash bottle used, drain plates on paper towels.

30

4. Use blocking buffer as diluent. Test sera, beginning at 1:50 dilution in the top row of the plate and dilute 2 fold/row down the plate (7X). Incubate at room temperature 1 hour. For subclone screening, a 1:1 dilution of culture supernatant in blocking buffer is used as starting material.

5. Repeat wash step #3.

6. Dilute secondary HRP-anti-hu IgG-Fc specific and/or HRP-anti-hu- κ reagent 1:2500-5000 in 1X PBST +5% chicken serum, add 100 μ L/well. Incubate 1 hour at room temperature.

7. Repeat wash step#3. (2X)

8. Develop plate using 10 mL citrate-phosphate buffer pH4.0,
80 μ L ABTS, 8 μ L H₂O₂ per plate.

5 9. Incubate 30 minutes to 1 hour at room temperature. Read
plate at OD_{415 nm-490 nm}. Consider twice above background titer limit.

In these assays, hybridomas 15H12 and 19D12 produced a positive ELISA
signal. These data demonstrate that the hybridomas produced antibodies which can
bind to soluble IGFR1.

10 Antigen positive hybridomas were then transferred to 24 well plates, and
eventually to tissue culture flasks. IGFR1 specific hybridomas were subcloned by
limiting dilution to assure monoclonality. Antigen positive hybridomas were preserved
at several stages in the development process by freezing cells in Origen DMSO
freeze medium (Fischer Scientific; Suwanee, GA).

15 **2.4.7.** Antibody isotypes were determined according to the following
procedure:

1. Coat plate overnight in refrigerator at 1 μ g/ml soluble IGFR1 in
1X PBS, 50 μ L/well. Empty plate.

2. Add 1X PBST + 5% chicken serum for 1 hour at room
temperature. (100 μ L/well). Empty plate.

20 3. Use blocking buffer as a diluent, add supernatant or purified
material to be tested in 1 well per secondary antibody to be tested-
50 μ L/well. Incubate for 90 minutes at room temperature. Empty plate.

25 4. Empty plate and wash manually with wash bottle (3X) or
plate washer (3X) using 1X PBST. If wash bottle used, drain plates on
paper towels.

30 5. Using blocking buffer as a diluent, add secondary antibodies:
HRP-anti-hu-gamma;
HRP-anti-hu kappa;
HRP-anti-human IgG1; or
HRP-anti-human IgG3
diluted 1:1000. Incubate for 45 minutes at room temperature. Empty
plate.

6. Repeat wash step #4 (3X).
7. Develop plate using 10 mL citrate-phosphate buffer pH4.0, 80 μ L ABTS, 8 μ L H₂O₂ per plate.
8. Incubate 30 minutes to 1 hour at room temperature. Read plate at OD_{415nm-490nm}.

The data from these assays is shown, below, in Table 4.

Table 4. Isotype ELISA results*.

	γ	κ	$\gamma 1$	$\gamma 3$			
	1	2	3	4	5	6	7
<u>clone</u>	1.903	1.003	0.064	0.813			
<u>15H12</u>							

*Each number represents the magnitude of the ELISA signal observed for each secondary antibody.

These data demonstrate that antibody 15H12 is an IgG3 κ antibody.

2.4.8. Hybridoma supernatants (1H3, 15H12 and 19D12) and MAB391 were also tested, in a fixed cell ELISA assay, for the ability to directly bind cells expressing IGFR1. In the assay, MCF-7 cells or HEK293 cells transfected with *IGFR1* DNA were used. The assays were performed as follows:

1. Add 50 μ g/well of a 20 μ g/mL solution of Poly-L-lysine in 1X PBS to each well of a 96 well plate and incubate for 30 minutes at room temperature or overnight at 4°C. Empty plate to remove Poly-L-lysine from the wells and allow to dry at room temperature until use.
2. Wash live cells three times with 1X PBS by centrifugation (1000 RPM/5 minutes). Adjust final cell concentration to 2 X 10⁶ cells per well in 1X PBS. Add 50 μ L per well of this cell suspension.
3. Spin cells 5 minutes at 2000 RPM. Empty buffer.
4. Add 50 μ L/well of 0.5% ice cold glutaraldehyde in 1X PBS. Let sit for 15 minutes at room temperature. Empty plate.
5. Add 1X PBST + 5% chicken serum and incubate for 1 hour at room temperature (100 μ L/well). Empty plate.

6. Wash plate gently using 1X PBST (2X). To avoid cell loss, this step should be done manually in a container avoiding any plate washers.

7. Using blocking buffer as a diluent, test culture supernatant by adding 100 μ g of a 1:1 dilution. Incubate 1 hour at room temperature.

8. Repeat step 6 (3X).

9. Dilute secondary HRP anti-hu IgG-Fc specific and/or HRP anti-hu- κ , reagent 1:2500-5000 in 1X PBST + 5% chicken serum, add 100 μ L/well. Incubate 1 hour at room temperature.

10. Repeat step 6 (3X).

11. Develop plate using 10 ml citrate-phosphate buffer pH4.0, 80 μ L, ABTS, 8 μ L H₂O₂ per plate.

12. Incubate 15-20 minutes at room temperature. Read plate at

OD_{415nm-490nm}.

The results from these assays demonstrated that hybridomas 1H3, 15H12 and 19D12 produced an immunoglobulin which binds to HEK293 cells expressing IGFR1 and that hybridomas 1H3, 15H12 and 19D12 produced an immunoglobulin which binds to MCF-7 cells which express endogenous IGFR1. Additionally, the results demonstrated that MAB391 bound to IGFR1 expressing HEK293 cells and to MCF-7 cells.

2.4.9. The ability of hybridoma supernatants (1H3, 15H12 and 19D12) to block binding of IGF1 to IGFR1 was evaluated by measuring 1) staining intensity of the supernatant on IGFR1 expressing HEK293 cells and on MCF7 cells and 2) the ability of the supernatants to block binding of IGF1-biotin to IGFR1 expressing cells. Initially, biotinylated IGF1 was titrated on IGFR1 expressing HEK293 cells in order to establish the proper concentration to evaluate blocking of IGF1 binding to its receptor by the antibodies of the present invention. This was done by the following procedure:

1. IGFR1 expressing HEK293 cells are harvested from a flask by slapping the flask to loosen the cells which were pipeted in to a conical tube. The cells are then centrifuged at 300 X g for 5 minutes to pellet the cells. The medium is then aspirated.

2. The cells are washed in 10-20 mL PBS containing 0.02% sodium azide and resuspended in the same buffer at approximately 2.5×10^6 cells/mL ($\pm 10^6$ cells). The cells are aliquoted, 200 μ L/well into a 96 well microtiter plate in the same buffer at 4°C. The cells are pelleted and the supernatant is aspirated.

3. The cells are stained by adding 50 μ L/well serially diluted IGF1-biotin in the same buffer, starting at a 1:5 dilution followed by 4-fold serial dilutions. The plate is tapped or gently vortexed to ensure an even suspension of cells are suspended. The cells are then incubated for 30 minutes at 4°C.

4. The cells are washed 3X by adding 150 μ L buffer for the first wash and then pelleted. The supernatant is aspirated and 200 μ L buffer is added. Again, the cells are pelleted and the supernatant is aspirated; this wash step is repeated once more. Streptavidin-PE (streptavidin-R-Phycoerythrin) is added and the cells are incubated for 30 minutes at 4°C.

5. The cells are washed once in PBS containing 2% FBS and 0.02% azide and resuspended in the same buffer except containing also 50 μ g/mL propidium iodide to exclude dead cells.

6. The cells are analyzed by FACS.

The blocking assays were performed as follows:

1. Harvest MCF7 cells or HEK293/IGFR1 cells from a tissue culture flask by slapping the flask sides to loosen the cells. Pipet the cells into a conical tube. Centrifuge the tube for 5 minutes at 300 X g to pellet the cells. Aspirate the medium.

2. Wash the cells in 10-20 mL PBS containing 2% FBS and 0.02% sodium azide (PFA), and resuspend in the same buffer at approximately 2.5×10^6 ($\pm 1 \times 10^6$). Aliquot 200 μ L/well into a 96 well microtiter plate in the same buffer at 4°C. Pellet the cells and aspirate the buffer.

3. Stain the cells with each IGFR1 hybridoma supernatant by adding 100 μ L/well, including a medium (negative) control, and MAB391 as a positive control. Tap the plate to ensure even suspension of the cells. Incubate 30-60 minutes at 4°C.

4. Wash the cells 3 times in PFA by adding 100 μ L buffer for the first wash, pellet, aspirate, resuspend in 200 μ L buffer, pellet, aspirate, resuspend again in 200 μ L buffer, divide each sample into two wells and pellet.

5 5. To one set of wells, add anti-human IgG-FITC diluted 1:100 in PFA (para-formaldehyde) to the supernatant stained samples and the medium control, and anti-mouse IgG-FITC at 1:200 to the MAB391 stained samples, again ensuring even dispersal of the cells (staining assay). Incubate for 30 minutes at 4°C.

10 6. To the second set of wells, add IGF1-biotin diluted 1:500 in PBS containing 0.02% azide (no FBS) and incubate for 30 minutes at 4°C. Wash the cells 3 times as described in step 4 (but without dividing the sample). Stain these cells by adding streptavidin-PE (streptavidin-R-Phycoerythrin) in PFA (blocking assay). Incubate for 30
15 minutes at 4°C.

7. Wash all the samples once in PFA, and resuspend in the same buffer except containing also 50 μ g/mL propidium iodide to exclude dead cells.

8. Analyze by FACS analysis.

20 The results from these blocking assays demonstrated that the supernatants from hybridomas 1H3, 15H12 and 19D12 block binding of biotinylated IGF1 to IGFR1, stain MCF7 cells which express endogenous IGFR1 and stain HEK293 cells expressing IGFR1.

25 **2.4.10.** The ability of purified antibodies 1H3 and 15H12 to block binding of biotinylated IGF1 to IGFR1 in an ELISA assay and of antibodies 1H3, 15H12 and 19D12 to block binding of biotinylated MAB391 to IGFR1 in an ELISA assay was also evaluated according to the following procedure:

30 1. Coat plate overnight in a refrigerator with 1 μ g/mL soluble IGFR1 in 1 X PBS-50 μ L/well.

2. Add 1 X PBST + 5% chicken serum for 1 hour at room temperature- 100 μ L/well. Empty plate.

3. Wash plate 3X with wash buffer (1X PBS + 0.05% tween-20). Slap plate dry.

4. 2 µg/mL 1H3, 15H12 or 19D12 or positive or negative control antibodies are diluted in blocking buffer across the plate. The plates are incubated at room temperature for 1 hour.

5. Wash plates 3X in wash buffer.

5 6. Biotin-IGF1 or Biotin-MAB391 is added-50µL/well- and incubated for 30 minutes at room temperature.

7. Wash plate 3X

8. Add 100µL/well of streptavidin labeled alkaline phosphatase or horse radish peroxidase, incubate for 30 minutes at room
10 temperature.

9. Wash plate 3X. Develop with appropriate reagent depending on the label used.

10. Read after 10-15 minutes.

MAB391 was biotinylated according to the following procedure:

15 1. Prepare MAB391 in PBS buffer (dialyze or use desalting column to remove unwanted buffers such as Tris or glycine).

2. Prepare a fresh stock solution of Sulfo-NHS-LC-biotin solution just before use. Add 2.0 mg of Sulfo-NHS-LC-biotin to 200 µL distilled water. Add this reagent to MAB391 at a 12-fold molar excess if
20 working with a 10 mg/mL solution of MAB391, or a 20-fold molar excess when working with a dilute preparation of MAB391 (2 mg/mL).

3. Calculation: mmoles MAB391 = mg protein/150,000
mmoles X 12 or 20 = mmoles biotin reagent to add
mmoles biotin to add X 556 = mg biotin reagent to add

25 For 1 mg/mL:

$$1/150000 = 6.6 \times 10^{-6}$$

$$20 \times 6.6 \times 10^{-6} \text{ mmoles} = 1/32 \times 10^{-4} \text{ NHS-LC-biotin}$$

$$1.32 \times 10^{-4} \times 556 = 0.073 \text{ mg sulfo NHS-LC-biotin}$$

From the stock NHS-LC-biotin solution, use 10 µL (100 µg) of
30 solution per mg IgG for 1 or 2 mg.

4. Incubate for 2 hours on ice or for 30 minutes at room temperature. Dialyze against PBS or use desalting column to remove unreacted biotin reagent. Store at 4°C in PBS 0.1% sodium azide.

In general, 3-5 biotins should be added to each IgG molecule labeled.

The results from these blocking assays demonstrated that antibodies 1H3 and 15H12 blocks binding of biotinylated IGF1 to sIGFR1 and that antibodies 1H3, 15H12 and 19D12 block biotinylated MAB391 binding to sIGFR1.

2.4.11. Binding between IGFR1 and the 1H3, 15H12 and 19D12 antibodies was evaluated in a BIAcore/surface plasmon resonance assay according to the following procedure:

1. IGFR1 is immobilized on a CM-5 chip by amine coupling, to a level of 350.4 response units on flow cells. The concentration of IGFR1 which is used to immobilize is 2.5 µg/mL in sodium acetate buffer and the protein is immobilized at pH 3.5.

2. Antibodies 1H3, 15H12 and 19D12 are purified from hybridoma supernatants over a Protein-A or Protein-G column and tested for purity by SDS-PAGE analysis (4%-12% Tris-Glycine).

3. The antibodies are made to flow over the IGFR1 surface prepared above.

4. The concentration range of antibodies used is 4, 2, 1, 0.5 and 0.25 µg/mL. A blank is also used for background substitution. Samples are prepared in HBS buffer.

5. Injection time (association phase) is 10 minutes, at a flow rate of 20 µL/minute, dissociation time (dissociation phase) is 1 hour at the same flow rate.

6. The assays are run at both 25°C and 37°C. All experiments are done in duplicate.

7. Data analysis is carried out using Bia-Evaluation software v.3.0.2 (Biacore, Inc; Piscataway, NJ).

8. All experiments are carried out using a Biacore 3000 surface plasmon resonance instrument (Biacore, Inc; Piscataway, NJ).

The results for these assays demonstrated that antibodies 15H12 and 19D12 associate with IGFR1 at 25°C and at 37°C and that antibody 1H3 associates with IGFR1 at 25°C. The data from these experiments were also used to calculate the affinity and rate constants of 1H3, 15H12 and 19D12 binding to IGFR1 (see Table 5, below).

Table 5. Affinity and rate constants of antibodies 1H3, 15H12 and 19D12 with IGFR1.

Temp.	Antibody	Sample size	Assoc. time (min.)	Dissoc. time (min.)	k_{on} (1/M's)	k_{off} (1/s)	K_D (M)	Half-life (min.) [#]
25°C	15H12	2	10	60	5.0×10^5	2.24×10^{-5}	4.48×10^{-11}	515.73
25°C	19D12	2	10	60	4.0×10^5	2.65×10^{-5}	5.92×10^{-11}	435.94
25°C	1H3	2	10	60	0.7×10^5	6.50×10^{-5}	86×10^{-11}	177.73
37°C	15H12	2	10	60	7.2×10^5	4.01×10^{-5}	5.57×10^{-11}	288.09
37°C	19D12	2	10	60	6.8×10^5	4.93×10^{-5}	7.22×10^{-11}	234.33

[#] Calculated as Half life = $\ln(2/k_{off})$

5

EXAMPLE 2: Cell Based Receptor Binding Assay.

A cell based receptor binding assay was used to determine if antibodies 1H3, 15H12 and 19D12 competed with IGF1 for binding to IGFR1.

In the assays, 96 well filter plates (1.2 μ m pore) were pre-wet with 0.5% bovine serum albumin (BSA)/PBS for 2 hours at 4°C. The buffer was then removed with a vacuum manifold. Various concentrations of 6X control or test antibody (1H3, 15H12 or 19D12) were added to the wells (25 μ L). The [¹²⁵I]-IGF-1 ligand was then added to the wells at a final concentration of 0.375 nM in BSA/PBS. Cells were harvested with cell dissociation solution, counted with trypan blue, and resuspended in 0.5% BSA/PBS to a cell number of 1-3 $\times 10^5$ /ml. One hundred μ L of cells (10,000-30,000) were added to each well. The plate was shaken at 4°C for 1 hour. The plate was then aspirated and washed three times with ice cold PBS using a vacuum manifold. The filters were punched out and counted on a gamma counter. Data were analyzed for competitive binding.

The results of these experiments indicated that 1H3, 15H12 and 19D12 were capable of competing with IGF-I for binding to IGFR1.

EXAMPLE 3: IGFR1 Autophosphorylation Assay.

The ability of 1H3, 15H12 and 19D12 to inhibit IGFR1 autophosphorylation was also determined.

Antibodies (1H3, 15H12 or 19D12) were added to cells bearing IGFR1 for various lengths of times. Cells were then stimulated with 10 ng/ml IGF-I for 5 min at

25

37°C. Cells were washed twice with cold PBS containing 0.1mM sodium vanadate and lysed in lysis buffer (50 mM HEPES, pH7.4, 150 mM NaCl, 10% glycerol, 1% Triton X-100, 1.5 mM MgCl₂, protease inhibitors and 2 mM sodium vanadate).

Lysates were incubated on ice for 30 min and then centrifuged at 13,000 RPM for 10 min at 4°C. Protein concentrations of the lysates were measured by a Coomassie colorimetric assay, and subjected to immunoprecipitation and Western blot analysis.

The results of these assays indicated that antibodies 1H3, 15H12 and 19D12 inhibited IGFR1 autophosphorylation with an IC₅₀ of 0.10 nM.

10 **EXAMPLE 4: Anchorage-Independent Growth (Soft Agar) Assay.**

The ability of an anti-IGFR1 antibodies 1H3, 15H12, 19D12 and MAB391 to inhibit anchorage-independent growth of various cells, including human breast cancer cell line MCF7, human colorectal cancer cell HT29 and human prostatic cancer cell DU145, was evaluated.

15 In these experiments, three milliliters of 0.6% agarose in complete MEM medium were added to each well of 6 well tissue culture plates and allowed to solidify (bottom layer). One hundred microliters of antibody 1H3, 15H12, 19D12 or MAB391 (discussed above), at various concentrations, was added to culture tubes. Cells were harvested. Aliquots of the cells (15,000 cells) were added to the culture tubes
20 containing the antibody and incubated at room temperature for 10-15 minutes. Three milliliters of a 0.35% agarose/complete minimal essential media (MEM) layer (top layer) were added to the antibody/cell mixture and then plated onto the solidified bottom layer. The top layer was allowed to solidify. The plates were then incubated for three weeks. MTT (3-(4,5-Dimethyl-2-Thiazolyl)-2,5-Diphenyl-2H-Tetrazolium
25 Bromide) was added to the wells and incubated for 1-2 hours. The plates were scanned and the colonies counted and analyzed using a customized colony counter application program.

The results of these experiments demonstrated that an anti-IGFR1 antibody can inhibit anchorage-independent growth of all three malignant cell lines tested.

30

EXAMPLE 5: Cloning of the Variable Regions of an Antibody from Hybridomas.

Nucleic acids encoding the 1H3, 15H12 and 19D12 variable regions were obtained from hybridomas according to the following procedure.

Messenger RNA (mRNA) from 2×10^8 hybridoma cells was prepared by using a Micro-Fast Track kit (Invitrogen; Carlsbad, CA). Cellular DNA (cDNA) encoding the variable region was prepared according the procedure described in "cDNA Cycle" kit (Invitrogen; Carlsbad, CA).

- 5 The antibody variable regions were PCR amplified using the cDNA as a template using 5'RACE (Clotech; Palo Alto, CA) technology. The following 3'primer sequence was used to amplify the heavy chain: 5'-TGCCAGGGGGAAGACCGATGG-3' (SEQ ID NO: 22) and following 3'primer sequence was used to amplify the light chain: 5'-CGGGAAGATGAAGACAGATG-3' (SEQ ID NO:23). Additionally, 5'-RACE
10 PCR primers (Clotech; Palo Alto, CA) were used in each amplification.

The PCR reaction mixture included 2.5 units of Pfu I polymerase in its appropriate buffer (Stratagene; La Jolla, CA), 0.2 mM of each dNTP, 750 nM of each 5' and 3' primer and cDNA template. Total reaction volume was 50 μ l. The following PCR cycling program was performed using a thermocycler:

- 15
- | | |
|-----|-----------------------------------|
| 1X | 94°C, 2 min. |
| 10X | 94°C, 45 sec. |
| | 65°C, 45 sec. Minus 1°C per cycle |
| | 72°C, 1 min. |
| 20 | 25X 94°C, 45 sec. |
| | 55°C, 45 sec. |
| | 72°C, 1 min. |
| | 1X 72°C, 15 min. |

- 25 The resulting PCR amplification product was inserted into the Zero Blunt TOPO PCR cloning vector (Invitrogen; Carlsbad, CA). The identity of the insert was verified by restriction enzyme analysis and then the nucleotide sequence of the insert was obtained by sequencing.

Example 6: Recombinant Expression of Antibody Chains

- 30 In this example, nucleic acids encoding various anti-IGFR1 antibody chains of the present invention were used to transfect a *dhfr*⁻ mammalian cell line (CHO-DXB11) wherein the chains were expressed. Transient transfections were carried out by cotransfection of the cell line with various combinations of one heavy (γ 1 or γ 4) and one light (κ) chain plasmid, selected from plasmids 1-11, listed below. Construction

of stable cell lines was performed by transfection by a single plasmid, either 12 or 13, listed below, as follows: The nucleic acids were located in a single plasmid and were operably linked to cytomegalovirus (CMV) promoters. The plasmids also contained *DHFR* cDNA operably linked to a mouse mammary tumor virus long terminal repeat (MMTV-LTR) which was used for plasmid amplification. The plasmid further included the hygromycin B gene operably linked to the *TK* promoter for selection in mammalian cells.

Below is a description of the promoter-expression cassette in the 13 plasmids which were constructed. The indicated plasmids (2-4 and 8-11) were deposited, under the Budapest Treaty, on _____ with the American Type Culture Collection (ATCC); 10801 University Boulevard; Manassas, Virginia 20110-2209 under the indicated name and accession number:

- (1) CMV promoter-15H12/19D12 HC (γ 4)
Insert Sequence: SEQ ID NO: 3;
- 15 (2) CMV promoter-15H12/19D12 HCA (γ 4)-
Deposit name: "15H12/19D12 HCA (γ 4)"
ATCC accession No.:
Insert Sequence: SEQ ID NO: 44;
- (3) CMV promoter-15H12/19D12 HCB (γ 4)-
20 Deposit name: "15H12/19D12 HCB (γ 4)"
ATCC accession No.:
Insert Sequence: SEQ ID NO: 111;
- (4) CMV promoter-15H12/19D12 HCA (γ 1)-
Deposit name: "15H12/19D12 HCA (γ 1)";
25 ATCC accession No.:
Insert Sequence: SEQ ID NO: 44;
- (5) CMV promoter-15H12/19D12 LC (κ)
Insert Sequence: SEQ ID NO: 1;
- (6) CMV promoter-15H12/19D12 LCA (κ)
30 Insert Sequence: SEQ ID NO: 40;
- (7) CMV promoter-15H12/19D12 LCB (κ)
Insert Sequence: SEQ ID NO: 42;
- (8) CMV promoter-15H12/19D12 LCC (κ)-

Deposit name: "15H12/19D12 LCC (κ)";

ATCC accession No.:

Insert Sequence: SEQ ID NO: 71;

(9) CMV promoter-15H12/19D12 LCD (κ)-

5 Deposit name: "15H12/19D12 LCD (κ)";

ATCC accession No.:

Insert Sequence: SEQ ID NO: 73;

(10) CMV promoter-15H12/19D12 LCE (κ)-

Deposit name: "15H12/19D12 LCE (κ)";

10 ATCC accession No.:

Insert Sequence: SEQ ID NO: 75;

(11) CMV promoter-15H12/19D12 LCF (κ)-

Deposit name: "15H12/19D12 LCF (κ)";

ATCC accession No.:

15 Insert Sequence: SEQ ID NO: 77;

(12) CMV promoter-15H12/19D12 HC (γ 4) and CMV promoter-15H12/19D12 LC (κ);

(13) CMV promoter-15H12/19D12 HCA (γ 1) and CMV promoter-15H12/19D12 LC (κ)

All restrictions on access to the plasmids deposited in ATCC will be removed upon grant of a patent.

20 The 3' end of each cassette was linked to a beta-globin poly A signal. The variable chains which were expressed were linked to the constant region indicated in parentheses (*i.e.*, γ 1, γ 4 or κ). Analysis of the transfected cell lines containing each plasmid indicated that the corresponding antibody chain polypeptides were expressed (amino acid sequences of the expression products not confirmed).

25 Each of the above-referenced plasmids constitutes part of the present invention. Further, the nucleic acid located within each expression cassette, along with the immunoglobulin variable region therein, along with the mature, processed version thereof (*i.e.*, lacking the signal sequence), particularly, SEQ ID NO: 44, mature HCA (nucleotides 58-411 of SEQ ID NO: 44), SEQ ID NO: 111, mature HCB
30 (nucleotides 58-411 of SEQ ID NO: 111), SEQ ID NO: 71, mature LCC (nucleotides 58-384 of SEQ ID NO: 71), SEQ ID NO: 73, mature LCD (nucleotides 58-384 of SEQ ID NO: 73), SEQ ID NO: 75, mature LCE (nucleotides 58-384 of SEQ ID NO: 75), SEQ ID NO: 77 or mature LCF (nucleotides 58-384 of SEQ ID NO: 77), optionally including an immunoglobulin constant region, along with any polypeptide encoded by

any of the foregoing nucleic acids, including mature or unprocessed chains, optionally including an immunoglobulin constant region, is a part of the present invention. Moreover, any antibody or antigen-binding fragment thereof comprising one of the encoded polypeptides is part of the present invention.

5

The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and the accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

10

Patents, patent applications, Genbank Accession Numbers and publications are cited throughout this application, the disclosures of which are incorporated herein by reference in their entireties.

15

WE CLAIM:

1. A binding composition that specifically binds to IGFR1 comprising a member selected from the group consisting of:
 - 5 a) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 8, CDR-L2 defined by SEQ ID NO: 9 and CDR-L3 defined by SEQ ID NO: 10;
 - b) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 31, CDR-L2 defined by SEQ ID NO: 32 and CDR-L3 defined by SEQ ID NO: 33;
 - 10 c) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 14 or SEQ ID NO: 17, CDR-H2 defined by SEQ ID NO: 15 and CDR-H3 defined by SEQ ID NO: 16; and
 - d) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 37 or SEQ ID NO: 70, CDR-H2 defined by SEQ ID NO: 38 and CDR-H3
 - 15 defined by SEQ ID NO: 39.

2. A binding composition of claim 1 that specifically binds to IGFR1 comprising a variable region selected from the group consisting of:
 - (a) amino acids 20-128 of SEQ ID NO: 2;
 - 20 (b) amino acids 21-130 of SEQ ID NO: 25;
 - (c) amino acids 20-128 of SEQ ID NO: 72;
 - (d) amino acids 20-128 of SEQ ID NO: 74.
 - (e) amino acids 20-137 of SEQ ID NO: 4;
 - (f) amino acids 20-140 of SEQ ID NO: 27;
 - 25 (g) amino acids 20-137 of SEQ ID NO: 45;
 - (h) amino acids 20-137 of SEQ ID NO: 112;
 - (i) amino acids 20-128 of SEQ ID NO: 76; and
 - (j) amino acids 20-128 of SEQ ID NO: 78.

- 30 3. A binding composition that specifically binds to IGFR1 comprising a member selected from the group consisting of:
 - (a) a light chain variable region comprising amino acids 20-128 of SEQ ID NO: 2 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 4;

- (b) a light chain variable region comprising amino acids 21-130 of SEQ ID NO: 25 and a heavy chain variable region comprising amino acids 20-140 of SEQ ID NO: 27;
 - (c) a light variable region comprising amino acids 20-128 of SEQ ID NO: 72 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 45;
 - 5 (d) a light variable region comprising amino acids 20-128 of SEQ ID NO: 74 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 45;
 - (e) a light variable region comprising amino acids 20-128 of SEQ ID NO: 76 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 45;
 - (f) a light variable region comprising amino acids 20-128 of SEQ ID NO: 78 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 45;
 - 10 (g) a light variable region comprising amino acids 20-128 of SEQ ID NO: 72 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 112;
 - (h) a light variable region comprising amino acids 20-128 of SEQ ID NO: 74 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 112;
 - 15 (i) a light variable region comprising amino acids 20-128 of SEQ ID NO: 76 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 112; and
 - (j) a light variable region comprising amino acids 20-128 of SEQ ID NO: 78 and a heavy chain variable region comprising amino acids 20-137 of SEQ ID NO: 112.
- 20 4. A pharmaceutical composition comprising a composition of claim 1 and a pharmaceutically acceptable carrier.
5. An isolated nucleic acid encoding a polypeptide selected from the group consisting of:
- 25 (a) amino acids 20-128 of SEQ ID NO: 2;
 - (b) amino acids 21-130 of SEQ ID NO: 25;
 - (c) amino acids 20-128 of SEQ ID NO: 72;
 - (d) amino acids 20-128 of SEQ ID NO: 74;
 - (e) amino acids 20-137 of SEQ ID NO: 4;
 - 30 (f) amino acids 20-140 of SEQ ID NO: 27;
 - (g) amino acids 20-137 of SEQ ID NO: 45;
 - (h) amino acids 20-137 of SEQ ID NO: 112;
 - (i) amino acids 20-128 of SEQ ID NO: 76; and
 - (j) amino acids 20-128 of SEQ ID NO: 78.

6. A nucleic acid of claim 5 selected from the group consisting of:
- (a) nucleotides 58-384 of SEQ ID NO: 1;
 - (b) nucleotides 61-390 of SEQ ID NO: 24;
 - 5 (c) nucleotides 58-384 of SEQ ID NO: 71;
 - (d) nucleotides 58-384 of SEQ ID NO: 73.
 - (e) nucleotides 58-411 of SEQ ID NO: 3;
 - (f) nucleotides 58-420 of SEQ ID NO: 26;
 - (g) nucleotides 58-411 of SEQ ID NO: 44;
 - 10 (h) nucleotides 58-411 of SEQ ID NO: 111;
 - (i) nucleotides 58-384 of SEQ ID NO: 75; and
 - (j) nucleotides 58-384 of SEQ ID NO: 77.
7. A recombinant vector comprising a nucleic acid of claim 5.
- 15 8. A host cell comprising a vector of claim 7.
9. A method for producing a polypeptide comprising culturing the host cell of claim 8 under conditions in which the polypeptide is produced.
- 20 10. A method for treating or preventing a medical condition in a subject, which medical condition is mediated by elevated expression or activity of Insulin-like Growth Factor Receptor-I, comprising administering a binding composition of claim 1 to the subject.
- 25 11. The method of claim 10 wherein the medical condition is selected from the group consisting of acromegaly, bladder cancer, Wilm's cancer, ovarian cancer, pancreatic cancer, benign prostatic hyperplasia, breast cancer, prostate cancer, bone cancer, lung cancer, colorectal cancer, cervical cancer, synovial sarcoma, diarrhea
- 30 associated with metastatic carcinoid, vasoactive intestinal peptide secreting tumors, gigantism, psoriasis, atherosclerosis, smooth muscle restenosis of blood vessels and inappropriate microvascular proliferation.

12. The method of claim 10 wherein the binding composition is administered to the subject by a parenteral route.

13. The method of claim 10 wherein the binding composition is administered to the subject in association with an additional, anti-cancer, therapeutic agent or anti-cancer, therapeutic procedure.

14. A method for treating or preventing a medical condition in a subject, which medical condition is mediated by elevated expression or activity of Insulin-like Growth Factor Receptor-I, comprising administering a binding composition that specifically binds to IGFR1 comprising a member selected from the group consisting of:

(a) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 8, CDR-L2 defined by SEQ ID NO: 9 and CDR-L3 defined by SEQ ID NO: 10;

(b) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 31, CDR-L2 defined by SEQ ID NO: 32 and CDR-L3 defined by SEQ ID NO: 33;

(c) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 14 or SEQ ID NO: 17, CDR-H2 defined by SEQ ID NO: 15 and CDR-H3 defined by SEQ ID NO: 16; and

(d) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 37 or SEQ ID NO: 70, CDR-H2 defined by SEQ ID NO: 38 and CDR-H3 defined by SEQ ID NO: 39;

to the subject.

15. The method of claim 14 wherein the medical condition is selected from the group consisting of acromegaly, bladder cancer, Wilm's cancer, ovarian cancer, pancreatic cancer, benign prostatic hyperplasia, breast cancer, prostate cancer, bone cancer, lung cancer, colorectal cancer, cervical cancer, synovial sarcoma, diarrhea associated with metastatic carcinoid, vasoactive intestinal peptide secreting tumors, gigantism, psoriasis, atherosclerosis, smooth muscle restenosis of blood vessels and inappropriate microvascular proliferation.

16. A method for producing a fully human, monoclonal antibody which specifically binds to IGFR1 comprising the steps of:

- (i) immunizing a transgenic non-human animal having a genome comprising a human heavy chain transgene and a human light chain transgene with an IGFR1 antigenic polypeptide, such that the antibody is produced by a B cell of the animal;
 - 5 (ii) isolating said B cell of the animal;
 - (iii) fusing the B cell with a myeloma cell to form an immortal hybridoma cell that secretes said antibody; and
 - (iv) isolating the antibody from the hybridoma cell.
- 10 17. The method of claim 16 wherein the antigenic polypeptide is amino acids 30-902 of SEQ ID NO: 19.
18. A binding composition which specifically binds to human IGFR1 comprising a property selected from the group consisting of:
- 15 (a) Binds to IGFR1 with a K_d of about 86×10^{-11} or less;
 - (b) Has an off rate (K_{off}) for IGFR1 of about 6.50×10^{-5} or smaller;
 - (c) Has an on rate (K_{on}) for IGFR1 of about 0.7×10^5 or greater;
 - (d) Competes with IGF1 for binding to IGFR1;
 - (e) Inhibits autophosphorylation of IGFR1; and
 - 20 (f) Inhibits anchorage-independent growth of a cell expressing IGFR1.
19. A binding composition of claim 18 comprising all of said properties.
20. A binding composition of claim 18 comprising a member selected from the group consisting of:
- 25 (a) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 8, CDR-L2 defined by SEQ ID NO: 9 and CDR-L3 defined by SEQ ID NO: 10;
 - (b) a light chain amino acid sequence which comprises CDR-L1 defined by SEQ ID NO: 31, CDR-L2 defined by SEQ ID NO: 32 and CDR-L3 defined by SEQ ID NO: 33;
 - 30 (c) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 14 or SEQ ID NO: 17, CDR-H2 defined by SEQ ID NO: 15 and CDR-H3 defined by SEQ ID NO: 16; and

- (d) a heavy chain amino acid sequence which comprises CDR-H1 defined by SEQ ID NO: 37 or SEQ ID NO: 70, CDR-H2 defined by SEQ ID NO: 38 and CDR-H3 defined by SEQ ID NO: 39.

SEQUENCE LISTING

<110> Schering Corp.

Wang, Yan

Pachter, Jonathan A

Hailey, Judith

Greenberg, Robert

Leonard, Presta

Brams, Peter

Feingersh, Diane

Williams, Denise

Srinivasan, Mohan

<120> NEUTRALIZING HUMAN ANTI-IGFR ANTIBODY

<130> OC01533-K-WI

<140>

<141>

<150> 60/383,459

<151> May 24, 2002

<150> 60/393,214

<151> July 2, 2002

<150> 60/436,254

<151> December 23, 2002

<160> 120

<170> PatentIn version 3.1

<210> 1

<211> 384

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> 1..384

<223>

<220>

<221> sig_peptide

<222> 1..57

<223>

<400> 1

atg tcg cca tca caa ctc att ggg ttt ctg ctg ctc tgg gtt cca gcc	48
Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala	
1 5 10 15	
tcc agg ggt gaa att gtg ctg act cag gtt cca gac ttt cag tct gtg	96
Ser Arg Gly Glu Ile Val Leu Thr Gln Val Pro Asp Phe Gln Ser Val	
20 25 30	
act cca aag gag aaa gtc acc atc acc tgc cgg gcc agt cag agc att	144
Thr Pro Lys Glu Lys Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile	
35 40 45	
ggt agt agc tta cac tgg tac cag cag aaa cca gat cag tct cca aag	192
Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys	
50 55 60	
ctc ctc atc aag tat gct tcc cag tcc ctc tca ggg gtc ccc tcg agg	240
Leu Leu Ile Lys Tyr Ala Ser Gln Ser Leu Ser Gly Val Pro Ser Arg	
65 70 75 80	
ttc agt ggc agt gga tct ggg aca gat ttc acc ctc acc atc aat agc	288
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser	
85 90 95	
ctg gaa gct gaa gat gct gca gcg tat tac tgt cat cag agt agt cgt	336
Leu Glu Ala Glu Asp Ala Ala Ala Tyr Tyr Cys His Gln Ser Ser Arg	

				100				105				110							
tta	cct	cac	act	ttc	ggc	gga	ggg	acc	aag	gtg	gag	atc	aaa	cga	act				
Leu	Pro	His	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys	Arg	Thr				
				115				120				125							
																384			

<210> 2

<211> 128

<212> PRT

<213> Homo sapiens

<400> 2

Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
1 5 10 15

Ser Arg Gly Glu Ile Val Leu Thr Gln Val Pro Asp Phe Gln Ser Val
20 25 30

Thr Pro Lys Glu Lys Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile
35 40 45

Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys
50 55 60

Leu Leu Ile Lys Tyr Ala Ser Gln Ser Leu Ser Gly Val Pro Ser Arg
65 70 75 80

Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser
85 90 95

Leu Glu Ala Glu Asp Ala Ala Ala Tyr Tyr Cys His Gln Ser Ser Arg
100 105 110

Leu Pro His Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr
115 120 125

<210> 3

<211> 411

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> 1..411

<223>

<220>

<221> sig_peptide

<222> 1..57

<223>

<400> 3

atg	gag	ttt	ggg	ctg	agc	tgg	gtt	ttc	ctt	gtt	gct	ata	tta	aaa	ggt	48
Met	Glu	Phe	Gly	Leu	Ser	Trp	Val	Phe	Leu	Val	Ala	Ile	Leu	Lys	Gly	
1			5					10						15		

gtc	cag	tgt	gag	gtt	cag	ctg	gtg	cag	tct	ggg	gga	ggc	ttg	gta	cat	96
Val	Gln	Cys	Glu	Val	Gln	Leu	Val	Gln	Ser	Gly	Gly	Gly	Leu	Val	His	
			20					25					30			

cct	ggg	ggg	tcc	ctg	aga	ctc	tcc	tgt	gca	gcc	tct	gga	ttc	acc	ttc	144
Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	
		35					40					45				

agt	agc	ttt	gct	atg	cac	tgg	gtt	cgc	cag	gct	cca	gga	aaa	ggt	ctg	192
Ser	Ser	Phe	Ala	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	
	50					55					60					

gag	tgg	ata	tca	gtt	att	gat	act	cgt	ggg	gcc	aca	tac	tat	gca	gac	240
Glu	Trp	Ile	Ser	Val	Ile	Asp	Thr	Arg	Gly	Ala	Thr	Tyr	Tyr	Ala	Asp	
65					70				75					80		

tcc	gtg	aag	ggc	cga	ttc	acc	atc	tcc	aga	gac	aat	gcc	aag	aac	tcc	288
Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Ser	
			85						90					95		

ttg	tat	ctt	caa	atg	aac	agc	ctg	aga	gcc	gag	gac	atg	gct	gtg	tat	336
Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Met	Ala	Val	Tyr	
			100					105					110			

tac	tgt	gca	aga	ctg	ggg	aac	ttc	tac	tac	ggg	atg	gac	gtc	tgg	ggc	384
Tyr	Cys	Ala	Arg	Leu	Gly	Asn	Phe	Tyr	Tyr	Gly	Met	Asp	Val	Trp	Gly	
		115					120					125				

caa	ggg	acc	acg	gtc	acc	gtc	tcc	tca								411
Gln	Gly	Thr	Thr	Val	Thr	Val	Ser	Ser								
		130				135										

<210> 4

<211> 137

<212> PRT

<213> Homo sapiens

<400> 4

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Lys Gly
 1 5 10 15

Val Gln Cys Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val His
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45

Ser Ser Phe Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

Glu Trp Ile Ser Val Ile Asp Thr Arg Gly Ala Thr Tyr Tyr Ala Asp
 65 70 75 80

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser
 85 90 95

Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr
 100 105 110

Tyr Cys Ala Arg Leu Gly Asn Phe Tyr Tyr Gly Met Asp Val Trp Gly
 115 120 125

Gln Gly Thr Thr Val Thr Val Ser Ser
 130 135

<210> 5

<211> 33

<212> DNA

<213> Homo sapiens

<400> 5

cgggccagtc agagcattgg tagtagctta cac

<210> 6

<211> 21

<212> DNA

<213> Homo sapiens

<400> 6

tatgcttccc agtccctctc a

21

<210> 7

<211> 27

<212> DNA

<213> Homo sapiens

<400> 7

catcagagta gtcgtttacc tcacact
24

<210> 8

<211> 11

<212> PRT

<213> Homo sapiens

<400> 8

Arg Ala Ser Gln Ser Ile Gly Ser Ser Leu His
1 5 10

<210> 9

<211> 7

<212> PRT

<213> Homo sapiens

<400> 9

Tyr Ala Ser Gln Ser Leu Ser
1 5

<210> 10

<211> 9

<212> PRT

<213> Homo sapiens

<400> 10

His Gln Ser Ser Arg Leu Pro His Thr
1 5

<210> 11

<211> 15

<212> DNA

<213> Homo sapiens

<400> 11

agctttgcta tgcac

15

<210> 12

<211> 48

<212> DNA

<213> Homo sapiens

<400> 12

gttattgata ctggtggtgc cacatactat gcagactccg tgaagggc

48

<210> 13

<211> 30

<212> DNA

<213> Homo sapiens

<400> 13

ctggggaact tctactacgg tatggacgtc

30

<210> 14

<211> 5

<212> PRT

<213> Homo sapiens

<400> 14

Ser Phe Ala Met His
1 5

<210> 15

<211> 16

<212> PRT

<213> Homo sapiens

<400> 15

Val Ile Asp Thr Arg Gly Ala Thr Tyr Tyr Ala Asp Ser Val Lys Gly
1 5 10 15

<210> 16

<211> 10

<212> PRT

<213> Homo sapiens

<400> 16

Leu Gly Asn Phe Tyr Tyr Gly Met Asp Val
1 5 10

<210> 17

<211> 10

<212> PRT

<213> Homo sapiens

<400> 17

Gly Phe Thr Phe Ser Ser Phe Ala Met His
1 5 10

<210> 18

<211> 30

<212> DNA

<213> Homo sapiens

<400> 18

ggattcacct tcagtagctt tgctatgcac
30

<210> 19

<211> 1367

<212> PRT

<213> Homo sapiens

<400> 19

Met Lys Ser Gly Ser Gly Gly Gly Ser Pro Thr Ser Leu Trp Gly Leu
1 5 10 15

Leu Phe Leu Ser Ala Ala Leu Ser Leu Trp Pro Thr Ser Gly Glu Ile
20 25 30

Cys Gly Pro Gly Ile Asp Ile Arg Asn Asp Tyr Gln Gln Leu Lys Arg
35 40 45

Leu Glu Asn Cys Thr Val Ile Glu Gly Tyr Leu His Ile Leu Leu Ile
50 55 60

Ser Lys Ala Glu Asp Tyr Arg Ser Tyr Arg Phe Pro Lys Leu Thr Val
65 70 75 80

Ile Thr Glu Tyr Leu Leu Leu Phe Arg Val Ala Gly Leu Glu Ser Leu
85 90 95

Gly Asp Leu Phe Pro Asn Leu Thr Val Ile Arg Gly Trp Lys Leu Phe
100 105 110

Tyr Asn Tyr Ala Leu Val Ile Phe Glu Met Thr Asn Leu Lys Asp Ile
115 120 125

Gly Leu Tyr Asn Leu Arg Asn Ile Thr Arg Gly Ala Ile Arg Ile Glu
130 135 140

Lys Asn Ala Asp Leu Cys Tyr Leu Ser Thr Val Asp Trp Ser Leu Ile
 145 150 155 160

Leu Asp Ala Val Ser Asn Asn Tyr Ile Val Gly Asn Lys Pro Pro Lys
 165 170 175

Glu Cys Gly Asp Leu Cys Pro Gly Thr Met Glu Glu Lys Pro Met Cys
 180 185 190

Glu Lys Thr Thr Ile Asn Asn Glu Tyr Asn Tyr Arg Cys Trp Thr Thr
 195 200 205

Asn Arg Cys Gln Lys Met Cys Pro Ser Thr Cys Gly Lys Arg Ala Cys
 210 215 220

Thr Glu Asn Asn Glu Cys Cys His Pro Glu Cys Leu Gly Ser Cys Ser
 225 230 235 240

Ala Pro Asp Asn Asp Thr Ala Cys Val Ala Cys Arg His Tyr Tyr Tyr
 245 250 255

Ala Gly Val Cys Val Pro Ala Cys Pro Pro Asn Thr Tyr Arg Phe Glu
 260 265 270

Gly Trp Arg Cys Val Asp Arg Asp Phe Cys Ala Asn Ile Leu Ser Ala
 275 280 285

Glu Ser Ser Asp Ser Glu Gly Phe Val Ile His Asp Gly Glu Cys Met
 290 295 300

Gln Glu Cys Pro Ser Gly Phe Ile Arg Asn Gly Ser Gln Ser Met Tyr
 305 310 315 320

Cys Ile Pro Cys Glu Gly Pro Cys Pro Lys Val Cys Glu Glu Glu Lys
 325 330 335

Lys Thr Lys Thr Ile Asp Ser Val Thr Ser Ala Gln Met Leu Gln Gly
 340 345 350

Cys Thr Ile Phe Lys Gly Asn Leu Leu Ile Asn Ile Arg Arg Gly Asn
 355 360 365

Asn Ile Ala Ser Glu Leu Glu Asn Phe Met Gly Leu Ile Glu Val Val
 370 375 380

Thr Gly Tyr Val Lys Ile Arg His Ser His Ala Leu Val Ser Leu Ser
385 390 395 400

Phe Leu Lys Asn Leu Arg Leu Ile Leu Gly Glu Glu Gln Leu Glu Gly
405 410 415

Asn Tyr Ser Phe Tyr Val Leu Asp Asn Gln Asn Leu Gln Gln Leu Trp
420 425 430

Asp Trp Asp His Arg Asn Leu Thr Ile Lys Ala Gly Lys Met Tyr Phe
435 440 445

Ala Phe Asn Pro Lys Leu Cys Val Ser Glu Ile Tyr Arg Met Glu Glu
450 455 460

Val Thr Gly Thr Lys Gly Arg Gln Ser Lys Gly Asp Ile Asn Thr Arg
465 470 475 480

Asn Asn Gly Glu Arg Ala Ser Cys Glu Ser Asp Val Leu His Phe Thr
485 490 495

Ser Thr Thr Thr Ser Lys Asn Arg Ile Ile Ile Thr Trp His Arg Tyr
500 505 510

Arg Pro Pro Asp Tyr Arg Asp Leu Ile Ser Phe Thr Val Tyr Tyr Lys
515 520 525

Glu Ala Pro Phe Lys Asn Val Thr Glu Tyr Asp Gly Gln Asp Ala Cys
530 535 540

Gly Ser Asn Ser Trp Asn Met Val Asp Val Asp Leu Pro Pro Asn Lys
545 550 555 560

Asp Val Glu Pro Gly Ile Leu Leu His Gly Leu Lys Pro Trp Thr Gln
565 570 575

Tyr Ala Val Tyr Val Lys Ala Val Thr Leu Thr Met Val Glu Asn Asp
580 585 590

His Ile Arg Gly Ala Lys Ser Glu Ile Leu Tyr Ile Arg Thr Asn Ala
595 600 605

Ser Val Pro Ser Ile Pro Leu Asp Val Leu Ser Ala Ser Asn Ser Ser
610 615 620

Ser Gln Leu Ile Val Lys Trp Asn Pro Pro Ser Leu Pro Asn Gly Asn
625 630 635 640

Leu Ser Tyr Tyr Ile Val Arg Trp Gln Arg Gln Pro Gln Asp Gly Tyr
645 650 655

Leu Tyr Arg His Asn Tyr Cys Ser Lys Asp Lys Ile Pro Ile Arg Lys
660 665 670

Tyr Ala Asp Gly Thr Ile Asp Ile Glu Glu Val Thr Glu Asn Pro Lys
675 680 685

Thr Glu Val Cys Gly Gly Glu Lys Gly Pro Cys Cys Ala Cys Pro Lys
690 695 700

Thr Glu Ala Glu Lys Gln Ala Glu Lys Glu Glu Ala Glu Tyr Arg Lys
705 710 715 720

Val Phe Glu Asn Phe Leu His Asn Ser Ile Phe Val Pro Arg Pro Glu
725 730 735

Arg Lys Arg Arg Asp Val Met Gln Val Ala Asn Thr Thr Met Ser Ser
740 745 750

Arg Ser Arg Asn Thr Thr Ala Ala Asp Thr Tyr Asn Ile Thr Asp Pro
755 760 765

Glu Glu Leu Glu Thr Glu Tyr Pro Phe Phe Glu Ser Arg Val Asp Asn
770 775 780

Lys Glu Arg Thr Val Ile Ser Asn Leu Arg Pro Phe Thr Leu Tyr Arg
785 790 795 800

Ile Asp Ile His Ser Cys Asn His Glu Ala Glu Lys Leu Gly Cys Ser
805 810 815

Ala Ser Asn Phe Val Phe Ala Arg Thr Met Pro Ala Glu Gly Ala Asp
820 825 830

Asp Ile Pro Gly Pro Val Thr Trp Glu Pro Arg Pro Glu Asn Ser Ile
835 840 845

Phe Leu Lys Trp Pro Glu Pro Glu Asn Pro Asn Gly Leu Ile Leu Met
850 855 860

Tyr Glu Ile Lys Tyr Gly Ser Gln Val Glu Asp Gln Arg Glu Cys Val

865		870		875		880
Ser Arg Gln Glu Tyr Arg Lys Tyr Gly Gly Ala Lys Leu Asn Arg Leu						
		885		890		895
Asn Pro Gly Asn Tyr Thr Ala Arg Ile Gln Ala Thr Ser Leu Ser Gly						
		900		905		910
Asn Gly Ser Trp Thr Asp Pro Val Phe Phe Tyr Val Gln Ala Lys Thr						
		915		920		925
Gly Tyr Glu Asn Phe Ile His Leu Ile Ile Ala Leu Pro Val Ala Val						
		930		935		940
Leu Leu Ile Val Gly Gly Leu Val Ile Met Leu Tyr Val Phe His Arg						
		945		950		955
						960
Lys Arg Asn Asn Ser Arg Leu Gly Asn Gly Val Leu Tyr Ala Ser Val						
		965		970		975
Asn Pro Glu Tyr Phe Ser Ala Ala Asp Val Tyr Val Pro Asp Glu Trp						
		980		985		990
Glu Val Ala Arg Glu Lys Ile Thr Met Ser Arg Glu Leu Gly Gln Gly						
		995		1000		1005
Ser Phe Gly Met Val Tyr Glu Gly Val Ala Lys Gly Val Val Lys						
		1010		1015		1020
Asp Glu Pro Glu Thr Arg Val Ala Ile Lys Thr Val Asn Glu Ala						
		1025		1030		1035
Ala Ser Met Arg Glu Arg Ile Glu Phe Leu Asn Glu Ala Ser Val						
		1040		1045		1050
Met Lys Glu Phe Asn Cys His His Val Val Arg Leu Leu Gly Val						
		1055		1060		1065
Val Ser Gln Gly Gln Pro Thr Leu Val Ile Met Glu Leu Met Thr						
		1070		1075		1080
Arg Gly Asp Leu Lys Ser Tyr Leu Arg Ser Leu Arg Pro Glu Met						
		1085		1090		1095
Glu Asn Asn Pro Val Leu Ala Pro Pro Ser Leu Ser Lys Met Ile						
		1100		1105		1110

Gln Met Ala Gly Glu Ile Ala Asp Gly Met Ala Tyr Leu Asn Ala
 1115 1120 1125

Asn Lys Phe Val His Arg Asp Leu Ala Ala Arg Asn Cys Met Val
 1130 1135 1140

Ala Glu Asp Phe Thr Val Lys Ile Gly Asp Phe Gly Met Thr Arg
 1145 1150 1155

Asp Ile Tyr Glu Thr Asp Tyr Tyr Arg Lys Gly Gly Lys Gly Leu
 1160 1165 1170

Leu Pro Val Arg Trp Met Ser Pro Glu Ser Leu Lys Asp Gly Val
 1175 1180 1185

Phe Thr Thr Tyr Ser Asp Val Trp Ser Phe Gly Val Val Leu Trp
 1190 1195 1200

Glu Ile Ala Thr Leu Ala Glu Gln Pro Tyr Gln Gly Leu Ser Asn
 1205 1210 1215

Glu Gln Val Leu Arg Phe Val Met Glu Gly Gly Leu Leu Asp Lys
 1220 1225 1230

Pro Asp Asn Cys Pro Asp Met Leu Phe Glu Leu Met Arg Met Cys
 1235 1240 1245

Trp Gln Tyr Asn Pro Lys Met Arg Pro Ser Phe Leu Glu Ile Ile
 1250 1255 1260

Ser Ser Ile Lys Glu Glu Met Glu Pro Gly Phe Arg Glu Val Ser
 1265 1270 1275

Phe Tyr Tyr Ser Glu Glu Asn Lys Leu Pro Glu Pro Glu Glu Leu
 1280 1285 1290

Asp Leu Glu Pro Glu Asn Met Glu Ser Val Pro Leu Asp Pro Ser
 1295 1300 1305

Ala Ser Ser Ser Ser Leu Pro Leu Pro Asp Arg His Ser Gly His
 1310 1315 1320

Lys Ala Glu Asn Gly Pro Gly Pro Gly Val Leu Val Leu Arg Ala
 1325 1330 1335

Ser Phe Asp Glu Arg Gln Pro Tyr Ala His Met Asn Gly Gly Arg
 1340 1345 1350

Lys Asn Glu Arg Ala Leu Pro Leu Pro Gln Ser Ser Thr Cys
 1355 1360 1365

<210> 20

<211> 195

<212> PRT

<213> Homo sapiens

<400> 20

Met Gly Lys Ile Ser Ser Leu Pro Thr Gln Leu Phe Lys Cys Cys Phe
 1 5 10 15

Cys Asp Phe Leu Lys Val Lys Met His Thr Met Ser Ser Ser His Leu
 20 25 30

Phe Tyr Leu Ala Leu Cys Leu Leu Thr Phe Thr Ser Ser Ala Thr Ala
 35 40 45

Gly Pro Glu Thr Leu Cys Gly Ala Glu Leu Val Asp Ala Leu Gln Phe
 50 55 60

Val Cys Gly Asp Arg Gly Phe Tyr Phe Asn Lys Pro Thr Gly Tyr Gly
 65 70 75 80

Ser Ser Ser Arg Arg Ala Pro Gln Thr Gly Ile Val Asp Glu Cys Cys
 85 90 95

Phe Arg Ser Cys Asp Leu Arg Arg Leu Glu Met Tyr Cys Ala Pro Leu
 100 105 110

Lys Pro Ala Lys Ser Ala Arg Ser Val Arg Ala Gln Arg His Thr Asp
 115 120 125

Met Pro Lys Thr Gln Lys Tyr Gln Pro Pro Ser Thr Asn Lys Asn Thr
 130 135 140

Lys Ser Gln Arg Arg Lys Gly Trp Pro Lys Thr His Pro Gly Gly Glu
 145 150 155 160

Gln Lys Glu Gly Thr Glu Ala Ser Leu Gln Ile Arg Gly Lys Lys Lys
 165 170 175

Glu Gln Arg Arg Glu Ile Gly Ser Arg Asn Ala Glu Cys Arg Gly Lys
 180 185 190

Lys Gly Lys
 195

<210> 21

<211> 180

<212> PRT

<213> Homo sapiens

<400> 21

Met Gly Ile Pro Met Gly Lys Ser Met Leu Val Leu Leu Thr Phe Leu
 1 5 10 15

Ala Phe Ala Ser Cys Cys Ile Ala Ala Tyr Arg Pro Ser Glu Thr Leu
 20 25 30

Cys Gly Gly Glu Leu Val Asp Thr Leu Gln Phe Val Cys Gly Asp Arg
 35 40 45

Gly Phe Tyr Phe Ser Arg Pro Ala Ser Arg Val Ser Arg Arg Ser Arg
 50 55 60

Gly Ile Val Glu Glu Cys Cys Phe Arg Ser Cys Asp Leu Ala Leu Leu
 65 70 75 80

Glu Thr Tyr Cys Ala Thr Pro Ala Lys Ser Glu Arg Asp Val Ser Thr
 85 90 95

Pro Pro Thr Val Leu Pro Asp Asn Phe Pro Arg Tyr Pro Val Gly Lys
 100 105 110

Phe Phe Gln Tyr Asp Thr Trp Lys Gln Ser Thr Gln Arg Leu Arg Arg
 115 120 125

Gly Leu Pro Ala Leu Leu Arg Ala Arg Arg Gly His Val Leu Ala Lys
 130 135 140

Glu Leu Glu Ala Phe Arg Glu Ala Lys Arg His Arg Pro Leu Ile Ala

145 150 155 160

Leu Pro Thr Gln Asp Pro Ala His Gly Gly Ala Pro Pro Glu Met Ala
 165 170 175

Ser Asn Arg Lys
 180

<210> 22

<211> 21

<212> DNA

<213> Artificial sequence

<220>

<223> PCR primer

<400> 22

tgccaggggg aagaccgatg g

21

<210> 23

<211> 20

<212> DNA

<213> Artificial sequence

<220>

<223> PCR primer

<400> 23

cggaagatg aagacagatg

20

<210> 24

<211> 390

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(390)

<223>

<220>

<221> sig_peptide

<222> (1)..(60)

<223>

<400> 24

atg	gaa	gcc	cca	gct	cag	ctt	ctc	ttc	ctc	ctg	cta	ctc	tgg	ctc	cca	48
Met	Glu	Ala	Pro	Ala	Gln	Leu	Leu	Phe	Leu	Leu	Leu	Leu	Trp	Leu	Pro	
1				5				10					15			

gat	acc	acc	gga	gaa	att	gtg	ttg	aca	cag	tct	cca	gcc	acc	ctg	tct	96
Asp	Thr	Thr	Gly	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	
			20					25					30			

ttg	tct	cca	ggg	gaa	aga	gcc	acc	ctc	tcc	tgc	agg	gcc	agt	cag	agt	144
Leu	Ser	Pro	Gly	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	
		35				40					45					

gtt	agc	agt	ttc	tta	gcc	tgg	tac	caa	cag	aaa	cct	ggc	cag	gct	ccc	192
Val	Ser	Ser	Phe	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	
	50					55					60					

agg	ctc	ctc	atc	tat	gat	gca	tcc	aac	agg	gcc	cct	ggc	atc	cca	gcc	240
Arg	Leu	Leu	Ile	Tyr	Asp	Ala	Ser	Asn	Arg	Ala	Pro	Gly	Ile	Pro	Ala	
65					70				75				80			

agg	ttc	agt	ggc	agt	ggg	tct	ggg	aca	gac	ttc	act	ctc	acc	atc	agc	288
Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	
			85					90					95			

agc	cta	gag	cct	gaa	gat	ttt	gca	gtt	tat	tac	tgt	cag	cag	cgt	agc	336
Ser	Leu	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Arg	Ser	
			100				105					110				

aac	tgg	cct	cgg	tgg	acg	ttc	ggc	caa	ggg	acc	aag	gtg	gaa	atc	aaa	384
Asn	Trp	Pro	Arg	Trp	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	
		115				120						125				

cga act

390

Arg Thr

130

<210> 25

<211> 130

<212> PRT

<213> Homo sapiens

<400> 25

Met Glu Ala Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro
1 5 10 15

Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser
20 25 30

Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser
35 40 45

Val Ser Ser Phe Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
50 55 60

Arg Leu Leu Ile Tyr Asp Ala Ser Asn Arg Ala Pro Gly Ile Pro Ala
65 70 75 80

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
85 90 95

Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Arg Ser
100 105 110

Asn Trp Pro Arg Trp Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
115 120 125

Arg Thr
130

<210> 26

<211> 420

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (420)

<223>

<220>

<221> sig_peptide

<222> (1)..(57)

<223>

<400> 26

atg	gag	ttt	ggg	ctg	agc	tgg	gtt	ttc	ctt	gtg	gct	ata	tta	aaa	ggt	48
Met	Glu	Phe	Gly	Leu	Ser	Trp	Val	Phe	Leu	Val	Ala	Ile	Leu	Lys	Gly	
1			5					10					15			

gtc	cag	tgt	gag	gtt	cag	ctg	gtg	cag	tct	ggg	gga	ggc	ttg	gta	ctt	96
Val	Gln	Cys	Glu	Val	Gln	Leu	Val	Gln	Ser	Gly	Gly	Gly	Leu	Val	Leu	
			20				25						30			

cct	ggg	ggg	tcc	ctg	aga	ctc	tcc	tgt	gca	ggc	tct	gga	ttc	acc	ttc	144
Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Gly	Ser	Gly	Phe	Thr	Phe	
			35				40					45				

agt	aac	tat	gct	atg	cac	tgg	att	cgc	cag	gct	cca	gga	aaa	ggt	ctg	192
Ser	Asn	Tyr	Ala	Met	His	Trp	Ile	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	
	50					55				60						

gag	tgg	gtg	tca	gct	att	ggt	gct	ggt	ggt	gac	acg	tac	tat	gca	gac	240
Glu	Trp	Val	Ser	Ala	Ile	Gly	Ala	Gly	Gly	Asp	Thr	Tyr	Tyr	Ala	Asp	
65					70					75					80	

tcc	gtg	aag	ggc	cga	ttc	acc	atc	tcc	aga	gac	aac	gcc	aag	gac	tcc	288
Ser	Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asp	Ser	
				85					90					95		

ttg	tat	ctt	caa	atg	aac	agc	ctg	aga	gcc	gag	gac	atg	gct	gtt	tat	336
Leu	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Met	Ala	Val	Tyr	
			100					105					110			

tac	tgt	gca	aga	ggc	cgg	cat	agg	aac	tgg	tac	tac	tac	aat	aag	gac	384
Tyr	Cys	Ala	Arg	Gly	Arg	His	Arg	Asn	Trp	Tyr	Tyr	Tyr	Asn	Lys	Asp	
		115				120						125				

tac	tgg	ggc	cag	gga	acc	ctg	gtc	acc	gtc	tcc	tca					
420																
Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser					
	130					135					140					

<210> 27

<211> 140

<212> PRT

<213> Homo sapiens

<400> 27

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Lys Gly
 1 5 10 15

Val Gln Cys Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Leu
 20 25 30

Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Gly Ser Gly Phe Thr Phe
 35 40 45

Ser Asn Tyr Ala Met His Trp Ile Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60

Glu Trp Val Ser Ala Ile Gly Ala Gly Gly Asp Thr Tyr Tyr Ala Asp
 65 70 75 80

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asp Ser
 85 90 95

Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Met Ala Val Tyr
 100 105 110

Tyr Cys Ala Arg Gly Arg His Arg Asn Trp Tyr Tyr Tyr Asn Lys Asp
 115 120 125

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 130 135 140

<210> 28

<211> 33

<212> DNA

<213> Homo sapiens

<400> 28

agggccagtc agagtgttag cagtttctta gcc

33

<210> 29

<211> 21

<212> DNA

<213> Homo sapiens

<400> 29
gatgcatcca acagggcccc t 21

<210> 30

<211> 30

<212> DNA

<213> Homo sapiens

<400> 30
cagcagcgta gcaactggcc tcggtggacg 30

<210> 31

<211> 11

<212> PRT

<213> Homo sapiens

<400> 31

Arg	Ala	Ser	Gln	Ser	Val	Ser	Ser	Phe	Leu	Ala
1				5					10	

<210> 32

<211> 7

<212> PRT

<213> Homo sapiens

<400> 32

Asp	Ala	Ser	Asn	Arg	Ala	Pro
1				5		

<210> 33

<211> 10

<212> PRT

<213> Homo sapiens

<400> 33

Gln Gln Arg Ser Asn Trp Pro Arg Trp Thr
1 5 10

<210> 34

<211> 30

<212> DNA

<213> Homo sapiens

<400> 34

ggattcacct tcagtaacta tgctatgcac
30

<210> 35

<211> 48

<212> DNA

<213> Homo sapiens

<400> 35

gctattggtg ctggtggtga cacgtactat gcagactccg tgaagggc
48

<210> 36

<211> 39

<212> DNA

<213> Homo sapiens

<400> 36

ggccggcata ggaactggta ctactacaat aaggactac
39

<210> 37

<211> 10

<212> PRT

<213> Homo sapiens

<400> 37

Gly Phe Thr Phe Ser Asn Tyr Ala Met His
 1 5 10

<210> 38

<211> 16

<212> PRT

<213> Homo sapiens

<400> 38

Ala Ile Gly Ala Gly Gly Asp Thr Tyr Tyr Ala Asp Ser Val Lys Gly
 1 5 10 15

<210> 39

<211> 13

<212> PRT

<213> Homo sapiens

<400> 39

Gly Arg His Arg Asn Trp Tyr Tyr Tyr Asn Lys Asp Tyr
 1 5 10

<210> 40

<211> 384

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (384)

<223>

<400> 40

atg tcg cca tca caa ctc att ggg ttt ctg ctg ctc tgg gtt cca gcc
 Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
 1 5 10 15

48

tcc agg ggt gaa att gtg ctg act cag agc cca gac tct ctg tct gtg 96
 Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val
 20 25 30

act cca ggc gag aga gtc acc atc acc tgc cgg gcc agt cag agc att 144
 Thr Pro Gly Glu Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile
 35 40 45

ggt agt agc tta cac tgg tac cag cag aaa cca ggt cag tct cca aag 192
 Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys
 50 55 60

ctt ctc atc tac tat gct tcc cag tcc ctc tca ggg gtc ccc tcg agg 240
 Leu Leu Ile Tyr Tyr Ala Ser Gln Ser Leu Ser Gly Val Pro Ser Arg
 65 70 75 80

ttc agt ggc agt gga tct ggg aca gat ttc acc ctc acc atc agt agc 288
 Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser
 85 90 95

ctc gag gct gaa gat ttc gca gtg tat tac tgt cat cag agt agt cgt 336
 Leu Glu Ala Glu Asp Phe Ala Val Tyr Tyr Cys His Gln Ser Ser Arg
 100 105 110

tta cct cac act ttc ggc caa ggg acc aag gtg gag atc aaa cgt acg 384
 Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 115 120 125

<210> 41

<211> 128

<212> PRT

<213> Homo sapiens

<400> 41

Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
1 5 10 15

Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val
 20 25 30

Thr Pro Gly Glu Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile
 35 40 45

Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys
 50 55 60

Leu Leu Ile Tyr Tyr Ala Ser Gln Ser Leu Ser Gly Val Pro Ser Arg
65 70 75 80

Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser
85 90 95

Leu Glu Ala Glu Asp Phe Ala Val Tyr Tyr Cys His Gln Ser Ser Arg
100 105 110

Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
115 120 125

<210> 42

<211> 384

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(384)

<223>

<400> 42
atg tcg cca tca caa ctc att ggg ttt ctg ctg ctc tgg gtt cca gcc 48
Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
1 5 10 15

tcc agg ggt gaa att gtg ctg act cag agc cca ggt acc ctg tct gtg 96
Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Val
20 25 30

tct cca ggc gag aga gcc acc ctc tcc tgc cgg gcc agt cag agc att 144
Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Ile
35 40 45

ggg agt agc tta cac tgg tac cag cag aaa cca ggt cag gct cca agg 192
Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg
50 55 60

ctt ctc atc tac tat gct tcc cag tcc ctc tca ggg atc ccc gat agg 240
Leu Leu Ile Tyr Tyr Ala Ser Gln Ser Leu Ser Gly Ile Pro Asp Arg
65 70 75 80

ttc agt ggc agt gga tct ggg aca gat ttc acc ctc acc atc agt aga 288
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg
85 90 95

ctg gag cct gaa gat ttc gca gtg tat tac tgt cat cag agt agt cgt 336
Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys His Gln Ser Ser Arg
100 105 110

tta cct cac act ttc ggc caa ggg acc aag gtg gag atc aaa cgt aca 384
 Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 115 120 125

<210> 43

<211> 128

<212> PRT

<213> Homo sapiens

<400> 43

Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
 1 5 10 15

Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Val
 20 25 30

Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Ile
 35 40 45

Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg
 50 55 60

Leu Leu Ile Tyr Tyr Ala Ser Gln Ser Leu Ser Gly Ile Pro Asp Arg
 65 70 75 80

Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg
 85 90 95

Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys His Gln Ser Ser Arg
 100 105 110

Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 115 120 125

<210> 44

<211> 411

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(411)

<223>

<400> 44

atg gag ttt ggg ctg agc tgg gtt ttc ctt gtt gct ata tta aaa ggt	48
Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Lys Gly	
1 5 10 15	
gtc cag tgt gag gtt cag ctg gtg cag tct ggg gga ggc ttg gta aag	96
Val Gln Cys Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Lys	
20 25 30	
cct ggg ggg tcc ctg aga ctc tcc tgt gca gcc tct gga ttc acc ttc	144
Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe	
35 40 45	
agt agc ttt gct atg cac tgg gtt cgc cag gct cca gga aaa ggt ctg	192
Ser Ser Phe Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu	
50 55 60	
gag tgg ata tca gtt att gat act cgt ggt gcc aca tac tat gca gac	240
Glu Trp Ile Ser Val Ile Asp Thr Arg Gly Ala Thr Tyr Tyr Ala Asp	
65 70 75 80	
tcc gtg aag ggc cga ttc acc atc tcc aga gac aat gcc aag aac tcc	288
Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser	
85 90 95	
ttg tat ctt caa atg aac agc ctg aga gcc gag gac act gct gtg tat	336
Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr	
100 105 110	
tac tgt gca aga ctg ggg aac ttc tac tac ggt atg gac gtc tgg ggc	384
Tyr Cys Ala Arg Leu Gly Asn Phe Tyr Tyr Gly Met Asp Val Trp Gly	
115 120 125	
caa ggg acc acg gtc acc gtc tcc tca	411
Gln Gly Thr Thr Val Thr Val Ser Ser	
130 135	

<210> 45

<211> 137

<212> PRT

<213> Homo sapiens

<400> 45

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Lys Gly

1					5					10					15				
Val	Gln	Cys	Glu	20	Val	Gln	Leu	Val	Gln	Ser	Gly	Gly	Gly	Leu	30	Val	Lys		
Pro	Gly	Gly	Ser	35	Leu	Arg	Leu	Ser	40	Cys	Ala	Ala	Ser	Gly	45	Phe	Thr		
Ser	Ser	Phe	Ala	50	Met	His	Trp	Val	55	Arg	Gln	Ala	Pro	Gly	60	Lys	Gly		
Glu	Trp	Ile	Ser	65	Val	Ile	Asp	Thr	70	Arg	Gly	Ala	Thr	Tyr	75	Tyr	Ala		
Ser	Val	Lys	Gly	85	Arg	Phe	Thr	Ile	Ser	90	Arg	Asp	Asn	Ala	Lys	95	Asn		
Leu	Tyr	Leu	Gln	100	Met	Asn	Ser	Leu	105	Arg	Ala	Glu	Asp	Thr	110	Val	Tyr		
Tyr	Cys	Ala	Arg	115	Leu	Gly	Asn	Phe	120	Tyr	Tyr	Gly	Met	Asp	125	Val	Trp		
Gln	Gly	Thr	Thr	130	Val	Thr	Val	Ser	135	Ser									

<210> 46

<211> 69

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (69)

<223>

```

<400> 46
gaa att gtg ctg act cag agc cca gac tct ctg tct gtg act cca ggc 48
Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val Thr Pro Gly
1 5 10 15
gag aga gtc acc atc acc tgc 69

```

Glu Arg Val Thr Ile Thr Cys
20

<210> 47

<211> 23

<212> PRT

<213> Homo sapiens

<400> 47

Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val Thr Pro Gly
1 5 10 15

Glu Arg Val Thr Ile Thr Cys
20

<210> 48

<211> 45

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(45)

<223>

<400> 48
tgg tac cag cag aaa cca ggt cag tct cca aag ctt ctc atc tac 45
Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr
1 5 10 15

<210> 49

<211> 15

<212> PRT

<213> Homo sapiens

<400> 49

Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr
 1 5 10 15

<210> 50

<211> 96

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(96)

<223>

<400> 50
 ggg gtc ccc tcg agg ttc agt ggc agt gga tct ggg aca gat ttc acc 48
 Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 1 5 10 15
 ctc acc atc agt agc ctc gag gct gaa gat ttc gca gtg tat tac tgt 96
 Leu Thr Ile Ser Ser Leu Glu Ala Glu Asp Phe Ala Val Tyr Tyr Cys
 20 25 30

<210> 51

<211> 32

<212> PRT

<213> Homo sapiens

<400> 51

Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 1 5 10 15
 Leu Thr Ile Ser Ser Leu Glu Ala Glu Asp Phe Ala Val Tyr Tyr Cys
 20 25 30

<210> 52

<211> 36

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (36)

<223>

<400> 52

ttc ggc caa ggg acc aag gtg gag atc aaa cgt acg
Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
1 5 10

36

<210> 53

<211> 12

<212> PRT

<213> Homo sapiens

<400> 53

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
1 5 10

<210> 54

<211> 69

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (69)

<223>

<400> 54

gaa att gtg ctg act cag agc cca ggt acc ctg tct gtg tct cca ggc
Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Val Ser Pro Gly
1 5 10 15

48

gag aga gcc acc ctc tcc tgc
 Glu Arg Ala Thr Leu Ser Cys
 20

69

<210> 55

<211> 23

<212> PRT

<213> Homo sapiens

<400> 55

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Val Ser Pro Gly
 1 5 10 15

Glu Arg Ala Thr Leu Ser Cys
 20

<210> 56

<211> 45

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(45)

<223>

<400> 56

tgg tac cag cag aaa cca ggt cag gct cca agg ctt ctc atc tac
 Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile Tyr
 1 5 10 15

45

<210> 57

<211> 15

<212> PRT

<213> Homo sapiens

<400> 57

Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile	Tyr
1				5					10					15

<210> 58

<211> 96

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(96)

<223>

<400> 58

ggg	atc	ccc	gat	agg	ttc	agt	ggc	agt	gga	tct	ggg	aca	gat	ttc	acc	48
Gly	Ile	Pro	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	
1				5					10					15		
ctc	acc	atc	agt	aga	ctg	gag	cct	gaa	gat	ttc	gca	gtg	tat	tac	tgt	96
Leu	Thr	Ile	Ser	Arg	Leu	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	
			20					25					30			

<210> 59

<211> 32

<212> PRT

<213> Homo sapiens

<400> 59

Gly	Ile	Pro	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr
1				5					10					15	
Leu	Thr	Ile	Ser	Arg	Leu	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys
			20					25					30		

<210> 60

<211> 36

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(39)

<223>

<400> 60

ttc ggc caa ggg acc aag gtg gag atc aaa cgt aca
Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
1 5 10

36

<210> 61

<211> 12

<212> PRT

<213> Homo sapiens

<400> 61

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
1 5 10

<210> 62

<211> 90

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(90)

<223>

<400> 62

gag gtt cag ctg gtg cag tct ggg gga ggc ttg gta aag cct ggg ggg
Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Lys Pro Gly Gly

48

1 5 10 15
 tcc ctg aga ctc tcc tgt gca gcc tct gga ttc acc ttc agt
 90
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
 20 25 30

<210> 63

<211> 30

<212> PRT

<213> Homo sapiens

<400> 63

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
 20 25 30

<210> 64

<211> 42

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(42)

<223>

<400> 64
 tgg gtt cgc cag gct cca gga aaa ggt ctg gag tgg ata tca
 Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ser
 1 5 10

42

<210> 65

<211> 14

<212> PRT

<213> Homo sapiens

<400> 65

Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ser
1 5 10

<210> 66

<211> 96

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(96)

<223>

<400> 66

cga ttc acc atc tcc aga gac aat gcc aag aac tcc ttg tat ctt caa 48
Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu Gln
1 5 10 15

atg aac agc ctg aga gcc gag gac act gct gtg tat tac tgt gca aga 96
Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
20 25 30

<210> 67

<211> 32

<212> PRT

<213> Homo sapiens

<400> 67

Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr Leu Gln
1 5 10 15

Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
20 25 30

<210> 68

<211> 33

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (33)

<223>

<400> 68

tgg ggc caa ggg acc acg gtc acc gtc tcc tca
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
1 5 10

33

<210> 69

<211> 11

<212> PRT

<213> Homo sapiens

<400> 69

Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
1 5 10

<210> 70

<211> 5

<212> PRT

<213> Homo sapiens

<400> 70

Asn Tyr Ala Met His
1 5

<210> 71

<211> 384

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(384)

<223>

```

<400> 71
atg tcg cca tca caa ctc att ggg ttt ctg ctg ctc tgg gtt cca gcc      48
Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
1                               5                               10          15

tcc agg ggt gaa att gtg ctg act cag agc cca gac tct ctg tct gtg      96
Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val
                20                               25          30

act cca ggc gag aga gtc acc atc acc tgc cgg gcc agt cag agc att      144
Thr Pro Gly Glu Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile
                35                               40          45

ggg agt agc tta cac tgg tac cag cag aaa cca ggt cag tct cca aag      192
Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys
                50                               55          60

ctt ctc atc aag tat gca tcc cag tcc ctc tca ggg gtc ccc tcg agg      240
Leu Leu Ile Lys Tyr Ala Ser Gln Ser Leu Ser Gly Val Pro Ser Arg
65                               70                               75          80

ttc agt ggc agt gga tct ggg aca gat ttc acc ctc acc atc agt agc      288
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser
                85                               90          95

ctc gag gct gaa gat gct gca gcg tat tac tgt cat cag agt agt cgt      336
Leu Glu Ala Glu Asp Ala Ala Tyr Tyr Cys His Gln Ser Ser Arg
                100                              105          110

tta cct cac act ttc ggc caa ggg acc aag gtg gag atc aaa cgt acg      384
Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
                115                              120          125

```

<210> 72

<211> 128

<212> PRT

<213> Homo sapiens

<400> 72

Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
 1 5 10 15

Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val
 20 25 30

Thr Pro Gly Glu Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile
 35 40 45

Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys
 50 55 60

Leu Leu Ile Lys Tyr Ala Ser Gln Ser Leu Ser Gly Val Pro Ser Arg
 65 70 75 80

Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser
 85 90 95

Leu Glu Ala Glu Asp Ala Ala Ala Tyr Tyr Cys His Gln Ser Ser Arg
 100 105 110

Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 115 120 125

<210> 73

<211> 384

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(384)

<223>

<400> 73

atg tcg cca tca caa ctc att ggg ttt ctg ctg ctc tgg gtt cca gcc
 Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
 1 5 10 15

48

tcc agg ggt gaa att gtg ctg act cag agc cca gac tct ctg tct gtg	96
Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val	
20 25 30	
act cca ggc gag aga gtc acc atc acc tgc cgg gcc agt cag agc att	144
Thr Pro Gly Glu Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile	
35 40 45	
ggg agt agc tta cac tgg tac cag cag aaa cca ggt cag tct cca aag	192
Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys	
50 55 60	
ctt ctc atc aag tat gca tcc cag tcc ctc tca ggg gtc ccc tcg agg	240
Leu Leu Ile Lys Tyr Ala Ser Gln Ser Leu Ser Gly Val Pro Ser Arg	
65 70 75 80	
ttc agt ggc agt gga tct ggg aca gat ttc acc ctc acc atc agt agc	288
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser	
85 90 95	
ctc gag gct gaa gat ttc gca gtg tat tac tgt cat cag agt agt cgt	336
Leu Glu Ala Glu Asp Phe Ala Val Tyr Tyr Cys His Gln Ser Ser Arg	
100 105 110	
tta cct cac act ttc ggc caa ggg acc aag gtg gag atc aaa cgt acg	384
Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr	
115 120 125	

<210> 74

<211> 128

<212> PRT

<213> Homo sapiens

<400> 74

Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala	
1 5 10 15	
Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val	
20 25 30	
Thr Pro Gly Glu Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile	
35 40 45	
Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys	
50 55 60	
Leu Leu Ile Lys Tyr Ala Ser Gln Ser Leu Ser Gly Val Pro Ser Arg	
65 70 75 80	

Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser
85 90 95

Leu Glu Ala Glu Asp Phe Ala Val Tyr Tyr Cys His Gln Ser Ser Arg
100 105 110

Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
115 120 125

<210> 75

<211> 384

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (384)

<223>

<400> 75
atg tcg cca tca caa ctc att ggg ttt ctg ctg ctc tgg gtt cca gcc 48
Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
1 5 10 15

tcc agg ggt gaa att gtg ctg act cag agc cca ggt acc ctg tct gtg 96
Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Val
20 25 30

tct cca ggc gag aga gcc acc ctc tcc tgc cgg gcc agt cag agc att 144
Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Ile
35 40 45

ggg agt agc tta cac tgg tac cag cag aaa cca ggt cag gct cca agg 192
Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg
50 55 60

ctt ctc atc aag tat gca tcc cag tcc ctc tca ggg atc ccc gat agg 240
Leu Leu Ile Lys Tyr Ala Ser Gln Ser Leu Ser Gly Ile Pro Asp Arg
65 70 75 80

ttc agt ggc agt gga tct ggg aca gat ttc acc ctc acc atc agt aga 288
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg
85 90 95

ctg gag cct gaa gat gct gca gcg tat tac tgt cat cag agt agt cgt 336
Leu Glu Pro Glu Asp Ala Ala Ala Tyr Tyr Cys His Gln Ser Ser Arg
100 105 110

tta cct cac act ttc ggc caa ggg acc aag gtg gag atc aaa cgt aca 384
 Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 115 120 125

<210> 76

<211> 128

<212> PRT

<213> Homo sapiens

<400> 76

Met Ser Pro Ser Gln Leu Ile Gly Phe Leu Leu Leu Trp Val Pro Ala
 1 5 10 15

Ser Arg Gly Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Val
 20 25 30

Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Ile
 35 40 45

Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg
 50 55 60

Leu Leu Ile Lys Tyr Ala Ser Gln Ser Leu Ser Gly Ile Pro Asp Arg
 65 70 75 80

Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg
 85 90 95

Leu Glu Pro Glu Asp Ala Ala Ala Tyr Tyr Cys His Gln Ser Ser Arg
 100 105 110

Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 115 120 125

<210> 77

<211> 384

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(384)

<223>

<400> 77

atg	tcg	cca	tca	caa	ctc	att	ggg	ttt	ctg	ctg	ctc	tgg	gtt	cca	gcc	48
Met	Ser	Pro	Ser	Gln	Leu	Ile	Gly	Phe	Leu	Leu	Leu	Trp	Val	Pro	Ala	
1				5					10					15		

tcc	agg	ggt	gaa	att	gtg	ctg	act	cag	agc	cca	ggt	acc	ctg	tct	gtg	96
Ser	Arg	Gly	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Gly	Thr	Leu	Ser	Val	
			20					25					30			

tct	cca	ggc	gag	aga	gcc	acc	ctc	tcc	tgc	cgg	gcc	agt	cag	agc	att	144
Ser	Pro	Gly	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Ile	
		35					40					45				

ggt	agt	agc	tta	cac	tgg	tac	cag	cag	aaa	cca	ggt	cag	gct	cca	agg	192
Gly	Ser	Ser	Leu	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	
	50					55					60					

ctt	ctc	atc	aag	tat	gca	tcc	cag	tcc	ctc	tca	ggg	atc	ccc	gat	agg	240
Leu	Leu	Ile	Lys	Tyr	Ala	Ser	Gln	Ser	Leu	Ser	Gly	Ile	Pro	Asp	Arg	
65					70					75				80		

ttc	agt	ggc	agt	gga	tct	ggg	aca	gat	ttc	acc	ctc	acc	atc	agt	aga	288
Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Arg	
				85					90					95		

ctg	gag	cct	gaa	gat	ttc	gca	gtg	tat	tac	tgt	cat	cag	agt	agt	cgt	336
Leu	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	His	Gln	Ser	Ser	Arg	
			100					105					110			

tta	cct	cac	act	ttc	ggc	caa	ggg	acc	aag	gtg	gag	atc	aaa	cgt	aca	384
Leu	Pro	His	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Arg	Thr	
		115					120					125				

<210> 78

<211> 128

<212> PRT

<213> Homo sapiens

<400> 78

Met	Ser	Pro	Ser	Gln	Leu	Ile	Gly	Phe	Leu	Leu	Leu	Trp	Val	Pro	Ala
1				5					10					15	

Ser	Arg	Gly	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Gly	Thr	Leu	Ser	Val
			20					25					30		

Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Ile
 35 40 45

Gly Ser Ser Leu His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg
 50 55 60

Leu Leu Ile Lys Tyr Ala Ser Gln Ser Leu Ser Gly Ile Pro Asp Arg
 65 70 75 80

Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg
 85 90 95

Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys His Gln Ser Ser Arg
 100 105 110

Leu Pro His Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 115 120 125

<210> 79

<211> 69

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(69)

<223>

<400> 79

gaa att gtg ctg act cag agc cca gac tct ctg tct gtg act cca ggc . 48
 Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val Thr Pro Gly
 1 5 10 15

gag aga gtc acc atc acc tgc . 69
 Glu Arg Val Thr Ile Thr Cys
 20

<210> 80

<211> 23

<212> PRT

<213> Homo sapiens

<400> 80

Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val Thr Pro Gly
1 5 10 15

Glu Arg Val Thr Ile Thr Cys
20

<210> 81

<211> 45

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(45)

<223>

<400> 81
tgg tac cag cag aaa cca ggt cag tct cca aag ctt ctc atc aag 45
Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Lys
1 5 10 15

<210> 82

<211> 15

<212> PRT

<213> Homo sapiens

<400> 82

Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Lys
1 5 10 15

<210> 83

<211> 96

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(96)

<223>

<400> 83

ggg gtc ccc tcg agg ttc agt ggc agt gga tct ggg aca gat ttc acc	48
Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr	
1 5 10 15	

ctc acc atc agt agc ctc gag gct gaa gat gct gca gcg tat tac tgt	96
Leu Thr Ile Ser Ser Leu Glu Ala Glu Asp Ala Ala Ala Tyr Tyr Cys	
20 25 30	

<210> 84

<211> 32

<212> PRT

<213> Homo sapiens

<400> 84

Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
1 5 10 15

Leu Thr Ile Ser Ser Leu Glu Ala Glu Asp Ala Ala Ala Tyr Tyr Cys
20 25 30

<210> 85

<211> 36

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (36)

<223>

<400> 85

ttc ggc caa ggg acc aag gtg gag atc aaa cgt acg
Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
1 5 10

36

<210> 86

<211> 12

<212> PRT

<213> Homo sapiens

<400> 86

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
1 5 10

<210> 87

<211> 69

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (69)

<223>

<400> 87

gaa att gtg ctg act cag agc cca gac tct ctg tct gtg act cca ggc
Glu Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ser Val Thr Pro Gly
1 5 10 15

48

gag aga gtc acc atc acc tgc
Glu Arg Val Thr Ile Thr Cys
20

69

<210> 88

<211> 23

<212> PRT

<213> Homo sapiens

<400> 88

Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Asp	Ser	Leu	Ser	Val	Thr	Pro	Gly
1				5					10					15	

Glu	Arg	Val	Thr	Ile	Thr	Cys
			20			

<210> 89

<211> 45

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(45)

<223>

<400> 89

tgg	tac	cag	cag	aaa	cca	ggg	cag	tct	cca	aag	ctt	ctc	atc	aag	
Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ser	Pro	Lys	Leu	Leu	Ile	Lys	45
1				5					10					15	

<210> 90

<211> 15

<212> PRT

<213> Homo sapiens

<400> 90

Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ser	Pro	Lys	Leu	Leu	Ile	Lys
1				5					10					15

<210> 91

<211> 96

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(96)

<223>

<400> 91

ggg gtc ccc tcg agg ttc agt ggc agt gga tct ggg aca gat ttc acc	48
Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr	
1 5 10 15	

ctc acc atc agt agc ctc gag gct gaa gat ttc gca gtg tat tac tgt	96
Leu Thr Ile Ser Ser Leu Glu Ala Glu Asp Phe Ala Val Tyr Tyr Cys	
20 25 30	

<210> 92

<211> 32

<212> PRT

<213> Homo sapiens

<400> 92

Gly Val Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
1 5 10 15

Leu Thr Ile Ser Ser Leu Glu Ala Glu Asp Phe Ala Val Tyr Tyr Cys
20 25 30

<210> 93

<211> 36

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(36)

<223>

<400> 93

ttc ggc caa ggg acc aag gtg gag atc aaa cgt acg
 Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 1 5 10

36

<210> 94

<211> 12

<212> PRT

<213> Homo sapiens

<400> 94

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 1 5 10

<210> 95

<211> 69

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(69)

<223>

<400> 95

gaa att gtg ctg act cag agc cca ggt acc ctg tct gtg tct cca ggc
 Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Val Ser Pro Gly
 1 5 10 15

48

gag aga gcc acc ctc tcc tgc
 Glu Arg Ala Thr Leu Ser Cys
 20

69

<210> 96

<211> 23

<212> PRT

<213> Homo sapiens

<400> 96

Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Gly	Thr	Leu	Ser	Val	Ser	Pro	Gly
1				5					10					15	

Glu	Arg	Ala	Thr	Leu	Ser	Cys
			20			

<210> 97

<211> 45

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(45)

<223>

tgg	tac	cag	cag	aaa	cca	ggt	cag	gct	cca	agg	ctt	ctc	atc	aag	
Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile	Lys	45
1				5					10					15	

<210> 98

<211> 15

<212> PRT

<213> Homo sapiens

<400> 98

Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	Arg	Leu	Leu	Ile	Lys
1				5					10					15

<220>

<221> CDS

<222> (1)..(36)

<223>

<400> 101

```

ttc ggc caa ggg acc aag gtg gag atc aaa cgt aca
Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
1           5           10

```

36

<210> 102

<211> 12

<212> PRT

<213> Homo sapiens

<400> 102

```

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
1           5           10

```

<210> 103

<211> 69

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(69)

<223>

<400> 103

```

gaa att gtg ctg act cag agc cca ggt acc ctg tct gtg tct cca ggc
Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Val Ser Pro Gly
1           5           10           15

```

48

```

gag aga gcc acc ctc tcc tgc
Glu Arg Ala Thr Leu Ser Cys

```

69

20

<210> 104

<211> 23

<212> PRT

<213> Homo sapiens

<400> 104

Glu Ile Val Leu Thr Gln Ser Pro Gly Thr Leu Ser Val Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys
20

<210> 105

<211> 45

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(45)

<223>

<400> 105
tgg tac cag cag aaa cca ggt cag gct cca agg ctt ctc atc aag 45
Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile Lys
1 5 10 15

<210> 106

<211> 15

<212> PRT

<213> Homo sapiens

<400> 106

Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile Lys
 1 5 10 15

<210> 107

<211> 96

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(96)

<223>

<400> 107

ggg atc ccc gat agg ttc agt ggc agt gga tct ggg aca gat ttc acc 48
 Gly Ile Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 1 5 10 15

ctc acc atc agt aga ctg gag cct gaa gat ttc gca gtg tat tac tgt 96
 Leu Thr Ile Ser Arg Leu Glu Pro Glu Asp Phe Ala Val Tyr Cys
 20 25 30

<210> 108

<211> 32

<212> PRT

<213> Homo sapiens

<400> 108

Gly Ile Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
 1 5 10 15

Leu Thr Ile Ser Arg Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys
 20 25 30

<210> 109

<211> 36

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(36)

<223>

<400> 109
 ttc ggc caa ggg acc aag gtg gag atc aaa cgt aca 36
 Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 1 5 10

<210> 110

<211> 12

<212> PRT

<213> Homo sapiens

<400> 110

Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
 1 5 10

<210> 111

<211> 411

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(411)

<223>

<400> 111
 atg gag ttt ggg ctg agc tgg gtt ttc ctt gtt gct ata tta aaa ggt 48
 Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Lys Gly
 1 5 10 15
 gtc cag tgt gag gtt cag ctg gtg cag tct ggg gga ggc ttg gta cag 96

Val Gln Cys Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Gln
 20 25 30
 ccc ggg ggg tcc ctg aga ctc tcc tgt gca gcc tct gga ttc acc ttc 144
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 agt agc ttt gct atg cac tgg gtt cgc cag gct cca gga aaa ggt ctg 192
 Ser Ser Phe Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 gag tgg ata tca gtt att gat act cgt ggt gcc aca tac tat gca gac 240
 Glu Trp Ile Ser Val Ile Asp Thr Arg Gly Ala Thr Tyr Tyr Ala Asp
 65 70 75 80
 tcc gtg aag ggc cga ttc acc atc tcc aga gac aat gcc aag aac tcc 288
 Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser
 85 90 95
 ttg tat ctt caa atg aac agc ctg aga gcc gag gac act gct gtg tat 336
 Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
 100 105 110
 tac tgt gca aga ctg ggg aac ttc tac tac ggt atg gac gtc tgg ggc 384
 Tyr Cys Ala Arg Leu Gly Asn Phe Tyr Tyr Gly Met Asp Val Trp Gly
 115 120 125
 caa ggg acc acg gtc acc gtc tcc tca 411
 Gln Gly Thr Thr Val Thr Val Ser Ser
 130 135

<210> 112

<211> 137

<212> PRT

<213> Homo sapiens

<400> 112

Met Glu Phe Gly Leu Ser Trp Val Phe Leu Val Ala Ile Leu Lys Gly
 1 5 10 15
 Val Gln Cys Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Gln
 20 25 30
 Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 35 40 45
 Ser Ser Phe Ala Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
 50 55 60
 Glu Trp Ile Ser Val Ile Asp Thr Arg Gly Ala Thr Tyr Tyr Ala Asp

65		70		75		80
Ser Val Lys Gly	Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser					
	85		90		95	
Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr						
	100		105		110	
Tyr Cys Ala Arg Leu Gly Asn Phe Tyr Tyr Gly Met Asp Val Trp Gly						
	115		120		125	
Gln Gly Thr Thr Val Thr Val Ser Ser						
	130		135			

<210> 113

<211> 75

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(75)

<223>

<400> 113	
gag gtt cag ctg gtg cag tct ggg gga ggc ttg gta cag cct ggg ggg	48
Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Gln Pro Gly Gly	
1 5 10 15	
tcc ctg aga ctc tcc tgt gca gcc tct	75
Ser Leu Arg Leu Ser Cys Ala Ala Ser	
20 25	

<210> 114

<211> 25

<212> PRT

<213> Homo sapiens

<400> 114

Glu Val Gln Leu Val Gln Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser
20 25

<210> 115

<211> 42

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(42)

<223>

<400> 115

tgg gtt cgc cag gct cca gga aaa ggt ctg gag tgg ata tca
Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ser
1 5 10

42

<210> 116

<211> 14

<212> PRT

<213> Homo sapiens

<400> 116

Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile Ser
1 5 10

<210> 117

<211> 96

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(96)

<223>

<400> 117

cga	ttc	acc	atc	tcc	aga	gac	aat	gcc	aag	aac	tcc	ttg	tat	ctt	caa	48
Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Ser	Leu	Tyr	Leu	Gln	
1				5				10						15		

atg	aac	agc	ctg	aga	gcc	gag	gac	act	gct	gtg	tat	tac	tgt	gca	aga	96
Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg	
			20				25						30			

<210> 118

<211> 32

<212> PRT

<213> Homo sapiens

<400> 118

Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Ser	Leu	Tyr	Leu	Gln
1				5				10						15	

Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	Ala	Arg
			20				25						30		

<210> 119

<211> 33

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(33)

<223>

<400> 119

tgg ggc caa ggg acc acg gtc acc gtc tcc tca
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
1 5 10

33

<210> 120

<211> 11

<212> PRT

<213> Homo sapiens

<400> 120

Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
1 5 10